



US007838708B2

(12) **United States Patent**  
**Sherman et al.**

(10) **Patent No.:** **US 7,838,708 B2**  
(45) **Date of Patent:** **Nov. 23, 2010**

(54) **HYDROCARBON CONVERSION PROCESS IMPROVEMENTS**

(75) Inventors: **Jeffrey H. Sherman**, Vero Beach, FL (US); **Eric W. McFarland**, Santa Barbara, CA (US); **Michael J. Weiss**, Santa Barbara, CA (US); **Ivan Marc Lorkovic**, Santa Barbara, CA (US); **Leroy E. Laverman**, Santa Barbara, CA (US); **Shouli Sun**, Santa Barbara, CA (US); **Dieter J. Schaefer**, Goleta, CA (US); **Galen D. Stucky**, Santa Barbara, CA (US); **Peter C. Ford**, Santa Barbara, CA (US); **Philip Grosso**, Auburn, CA (US); **Ashley W. Breed**, Goleta, CA (US); **Michael F. Doherty**, Santa Barbara, CA (US)

(73) Assignees: **GRT, Inc.**, Santa Barbara, CA (US); **The Regents of the University of California**, Oakland, CA (US)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/692,831**

(22) Filed: **Jan. 25, 2010**

(65) **Prior Publication Data**

US 2010/0121119 A1 May 13, 2010

**Related U.S. Application Data**

(63) Continuation of application No. 11/098,997, filed on Apr. 5, 2005, now abandoned, which is a continuation-in-part of application No. 10/430,240, filed on Aug. 19, 2003, now Pat. No. 7,161,050, which is a continuation-in-part of application No. 10/365,346, filed on Feb. 12, 2003, now abandoned, which is a continuation of application No. 10/298,440, filed on Nov. 19, 2002, now abandoned, which is a continuation-in-part of application No. 10/208,068, filed on Jul. 29, 2002, now abandoned, which is a continuation-in-part of application No. 10/054,004, filed on Jan. 24, 2002, now Pat. No. 6,486,368, which is a continuation-in-part of application No. 09/951,739, filed on Sep. 11, 2001, now Pat. No. 6,465,696, which is a continuation-in-part of application No. 09/886,078, filed on Jun. 20, 2001, now Pat. No. 6,472,572.

(51) **Int. Cl.**

**C07C 29/48** (2006.01)  
**C07C 29/58** (2006.01)

(52) **U.S. Cl.** ..... **568/893; 568/910; 568/910.5**

(58) **Field of Classification Search** ..... **568/893, 568/910, 910.5**

See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

2,168,260 A 8/1939 Heisel et al.

2,246,082 A	6/1941	Vaughan et al.
2,488,083 A	11/1949	Gorin et al.
2,677,598 A	5/1954	Crummett et al.
2,941,014 A	6/1960	Rothweiler et al.
3,076,784 A	2/1963	Huermann et al.
3,172,915 A	3/1965	Borkowski et al.
3,246,043 A	4/1966	Rosset et al.
3,273,964 A	9/1966	Rosset
3,294,846 A	12/1966	Livak et al.
3,310,380 A	3/1967	Lester
3,346,340 A	10/1967	Louvar et al.
3,353,916 A	11/1967	Lester
3,353,919 A	11/1967	Stockman
3,496,242 A	2/1970	Berkowitz et al.
3,562,321 A	2/1971	Borkowski et al.
3,598,876 A	8/1971	Bloch
3,657,367 A	4/1972	Blake et al.
3,670,037 A	6/1972	Dugan
3,673,264 A	6/1972	Kuhn
3,679,758 A	7/1972	Schneider
3,702,886 A	11/1972	Argauer et al.
3,705,196 A	12/1972	Turner
3,799,997 A	3/1974	Schmerling

(Continued)

**FOREIGN PATENT DOCUMENTS**

BR	0210054	8/2004
CA	1099656	4/1981
CA	1101441	5/1981
CA	1202610	4/1986
CA	2447761 A1	11/2002
CA	2471295 A1	7/2003
CA	2542857	5/2005
CA	2236126	8/2006
CA	2203115	9/2006
CA	2510093	12/2006

(Continued)

**OTHER PUBLICATIONS**

U.S. Appl. No. 60/487,364, filed Jul. 15, 2003, Lorkovic et al.  
U.S. Appl. No. 60/559,844, filed Apr. 6, 2004, Sherman et al.  
U.S. Appl. No. 60/765,115, filed Feb. 3, 2006, Gadewar et al.  
Abstract of JP2007045756, Hydrogenation method using diaphragm type hydrogenation catalyst, hydrogenation reaction apparatus and diaphragm type hydrogenation catalyst, Publication date: Feb. 22, 2007, Inventor: Shuji et al., esp@cenet database—worldwide.

(Continued)

*Primary Examiner*—Elvis O Price

(74) *Attorney, Agent, or Firm*—Baker Botts L.L.P.

(57) **ABSTRACT**

Improvements in previously disclosed methods of and apparatuses for converting alkanes, alkenes, and aromatics to olefins, alcohols, ethers, and aldehydes includes: safety improvements, use of alternative feedstocks, process simplification, improvements to the halogenation step, improvements to the reproporation step, improvements to the solid oxide reaction, improvements to solid oxide regeneration, improvements in separations, maintenance, start-up, shut-down, and materials of construction.

**20 Claims, No Drawings**

# US 7,838,708 B2

Page 2

---

## U.S. PATENT DOCUMENTS

3,865,886 A	2/1975	Schindler et al.	4,371,716 A	2/1983	Paxson et al.
3,876,715 A	4/1975	McNulty et al.	4,373,109 A	2/1983	Olah
3,879,473 A	4/1975	Stapp	4,376,019 A	3/1983	Gamlen et al.
3,879,480 A	4/1975	Riegel et al.	4,380,682 A	4/1983	Leitert et al.
3,883,651 A	5/1975	Woitun et al.	4,384,159 A	5/1983	Diesen
3,886,287 A	5/1975	Kobayashi et al.	4,389,391 A	6/1983	Dunn
3,894,103 A	7/1975	Chang et al.	4,410,714 A	10/1983	Apanel
3,894,104 A	7/1975	Chang et al.	4,412,086 A	10/1983	Beard, Jr. et al.
3,894,105 A	7/1975	Chang et al.	4,418,236 A	11/1983	Cornelius et al.
3,894,107 A	7/1975	Butter et al.	4,431,856 A	2/1984	Daviduk et al.
3,907,917 A	9/1975	Forth	4,433,189 A	2/1984	Young
3,919,336 A	11/1975	Kurtz	4,433,192 A	2/1984	Olah
3,920,764 A	11/1975	Riegel et al.	4,439,409 A	3/1984	Puppe et al.
3,923,913 A	12/1975	Antonini et al.	4,440,871 A	4/1984	Lok et al.
3,928,483 A	12/1975	Chang et al.	4,443,620 A	4/1984	Gelbein et al.
3,965,205 A	6/1976	Garwood et al.	4,462,814 A	7/1984	Holmes et al.
3,974,062 A	8/1976	Owen et al.	4,465,884 A	8/1984	Degnan et al.
3,987,119 A	10/1976	Kurtz et al.	4,467,130 A	8/1984	Olah
3,992,466 A	11/1976	Plank et al.	4,467,133 A	8/1984	Chang et al.
4,006,169 A	2/1977	Anderson et al.	4,489,210 A	12/1984	Judat et al.
4,011,278 A	3/1977	Plank et al.	4,489,211 A	12/1984	Ogura et al.
4,025,571 A	5/1977	Lago	4,492,657 A	1/1985	Heiss
4,025,572 A	5/1977	Lago	4,496,752 A	1/1985	Gelbein et al.
4,025,575 A	5/1977	Chang et al.	4,497,967 A	2/1985	Wan
4,025,576 A	5/1977	Chang et al.	4,499,314 A	2/1985	Seddon et al.
4,035,285 A	7/1977	Owen et al.	4,506,105 A	3/1985	Kaufhold
4,035,430 A	7/1977	Dwyer et al.	4,509,955 A	4/1985	Hayashi
4,039,600 A	8/1977	Chang	4,513,092 A	4/1985	Chu et al.
4,044,061 A	8/1977	Chang et al.	4,513,164 A	4/1985	Olah
4,046,825 A	9/1977	Owen et al.	4,523,040 A	6/1985	Olah
4,049,734 A	9/1977	Garwood et al.	4,524,227 A	6/1985	Fowles et al.
4,052,471 A	10/1977	Pearsall	4,524,228 A	6/1985	Fowles et al.
4,052,472 A	10/1977	Given et al.	4,524,231 A	6/1985	Fowles et al.
4,058,576 A	11/1977	Chang et al.	4,538,014 A	8/1985	Miale et al.
4,060,568 A	11/1977	Rodewald	4,538,015 A	8/1985	Miale et al.
4,071,753 A	1/1978	Fulenwider et al.	4,540,826 A	9/1985	Banasiak et al.
4,072,733 A	2/1978	Hargis et al.	4,543,434 A	9/1985	Chang
4,087,475 A	5/1978	Jordan	4,544,781 A	10/1985	Chao et al.
4,088,706 A	5/1978	Kaeding	4,547,612 A	10/1985	Tabak
4,092,368 A	5/1978	Smith	4,550,217 A	10/1985	Graziani et al.
4,110,180 A	8/1978	Nidola et al.	4,550,218 A	10/1985	Chu
4,117,251 A	9/1978	Kaufhold et al.	4,568,660 A	2/1986	Klosiewicz
4,129,604 A	12/1978	Tsao	4,579,977 A	4/1986	Drake
4,133,838 A	1/1979	Pearson	4,579,992 A	4/1986	Kaufhold et al.
4,133,966 A	1/1979	Pretzer et al.	4,579,996 A	4/1986	Font Freide et al.
4,138,440 A	2/1979	Chang et al.	4,587,375 A	5/1986	Debras et al.
4,156,698 A	5/1979	Dwyer et al.	4,588,835 A	5/1986	Torii et al.
4,169,862 A	10/1979	Eden	4,590,310 A	5/1986	Townsend et al.
4,172,099 A	10/1979	Severino	4,599,474 A	7/1986	Devries et al.
4,187,255 A	2/1980	Dodd	4,605,796 A	8/1986	Isogai et al.
4,194,990 A	3/1980	Pieters et al.	4,605,803 A	8/1986	Chang et al.
4,197,420 A	4/1980	Ferraris et al.	4,621,161 A	11/1986	Shihabi
4,219,680 A	8/1980	Konig et al.	4,621,164 A	11/1986	Chang et al.
4,249,031 A	2/1981	Drent et al.	4,633,027 A	12/1986	Owen et al.
4,270,929 A	6/1981	Dang Vu et al.	4,634,800 A	1/1987	Withers, Jr. et al.
4,272,338 A	6/1981	Lynch et al.	4,642,403 A	2/1987	Hyde et al.
4,282,159 A	8/1981	Davidson et al.	4,642,404 A	2/1987	Shihabi
4,300,005 A	11/1981	Li	4,652,688 A	3/1987	Brophy et al.
4,300,009 A	11/1981	Haag et al.	4,654,449 A	3/1987	Chang et al.
4,301,253 A	11/1981	Warren	4,655,893 A	4/1987	Beale
4,302,619 A	11/1981	Gross et al.	4,658,073 A	4/1987	Tabak
4,307,261 A	12/1981	Beard, Jr. et al.	4,658,077 A	4/1987	Kolts et al.
4,308,403 A	12/1981	Knifton	4,665,259 A	5/1987	Brazdil et al.
4,311,865 A	1/1982	Chen et al.	4,665,267 A	5/1987	Barri
4,317,800 A	3/1982	Sloterdijk et al.	4,665,270 A	5/1987	Brophy et al.
4,317,934 A	3/1982	Seemuth	4,675,410 A	6/1987	Feitler et al.
4,317,943 A	3/1982	Knifton	4,690,903 A	9/1987	Chen et al.
4,320,241 A	3/1982	Frankiewicz	4,695,663 A	9/1987	Hall et al.
4,333,852 A	6/1982	Warren	4,696,985 A	9/1987	Martin
4,347,391 A	8/1982	Campbell	4,704,488 A	11/1987	Devries et al.
4,350,511 A	9/1982	Holmes et al.	4,704,493 A	11/1987	Devries et al.
			4,709,108 A	11/1987	Devries et al.

# US 7,838,708 B2

Page 3

---

4,720,600 A	1/1988	Beech, Jr. et al.	5,082,473 A	1/1992	Keefer
4,720,602 A	1/1988	Chu	5,082,816 A	1/1992	Teller et al.
4,724,275 A	2/1988	Hinnenkamp et al.	5,085,674 A	2/1992	Leavitt
4,735,747 A	4/1988	Ollivier et al.	5,087,779 A	2/1992	Nubel et al.
4,737,594 A	4/1988	Olah	5,087,786 A	2/1992	Nubel et al.
4,748,013 A	5/1988	Saito et al.	5,087,787 A	2/1992	Kimble et al.
4,769,504 A	9/1988	Noceti et al.	5,093,542 A	3/1992	Gaffney
4,774,216 A	9/1988	Kolts et al.	5,096,469 A	3/1992	Keefer
4,775,462 A	10/1988	Imai et al.	5,097,083 A	3/1992	Stauffer
4,777,321 A	10/1988	Harandi et al.	5,099,084 A	3/1992	Stauffer
4,781,733 A	11/1988	Babcock et al.	5,105,045 A	4/1992	Kimble et al.
4,783,566 A	11/1988	Kocal et al.	5,105,046 A	4/1992	Washecheck
4,788,369 A	11/1988	Marsh et al.	5,107,032 A	4/1992	Erb et al.
4,788,377 A	11/1988	Chang et al.	5,107,051 A	4/1992	Pannell
4,792,642 A	12/1988	Rule et al.	5,107,061 A	4/1992	Ou et al.
4,795,732 A	1/1989	Barri	5,108,579 A	4/1992	Casci
4,795,737 A	1/1989	Rule et al.	5,118,899 A	6/1992	Kimble et al.
4,795,843 A	1/1989	Imai et al.	5,120,332 A	6/1992	Wells
4,795,848 A	1/1989	Teller et al.	5,132,343 A	7/1992	Zwecker et al.
4,804,797 A	2/1989	Minet et al.	5,138,112 A	8/1992	Gosling et al.
4,804,800 A	2/1989	Bortinger et al.	5,139,991 A	8/1992	Taylor et al.
4,808,763 A	2/1989	Shum	5,146,027 A	9/1992	Gaffney
4,814,527 A	3/1989	Diesen	5,157,189 A	10/1992	Karra
4,814,532 A	3/1989	Yoshida et al.	5,160,502 A	11/1992	Kimble et al.
4,814,535 A	3/1989	Yurchak	5,166,452 A	11/1992	Gradl et al.
4,814,536 A	3/1989	Yuchak	5,175,382 A	12/1992	Hebgen et al.
4,849,562 A	7/1989	Buhs et al.	5,178,748 A	1/1993	Casci et al.
4,849,573 A	7/1989	Kaefing	5,185,479 A	2/1993	Stauffer
4,851,602 A	7/1989	Harandi et al.	5,188,725 A	2/1993	Harandi
4,851,606 A	7/1989	Ragonese et al.	5,191,142 A	3/1993	Marshall et al.
4,886,925 A	12/1989	Harandi	5,194,244 A	3/1993	Brownscombe et al.
4,886,932 A	12/1989	Leyshon	5,202,506 A	4/1993	Kirchner et al.
4,891,463 A	1/1990	Chu	5,202,511 A	4/1993	Salinas, III et al.
4,895,995 A	1/1990	James, Jr. et al.	5,210,357 A	5/1993	Kolts et al.
4,899,000 A	2/1990	Stauffer	5,215,648 A	6/1993	Zones et al.
4,899,001 A	2/1990	Kalnes et al.	5,223,471 A	6/1993	Washecheck
4,899,002 A	2/1990	Harandi et al.	5,228,888 A	7/1993	Gmelin et al.
4,902,842 A	2/1990	Kalnes et al.	5,233,113 A	8/1993	Periana et al.
4,925,995 A	5/1990	Robschlag	5,237,115 A	8/1993	Makovec et al.
4,929,781 A	5/1990	James, Jr. et al.	5,243,098 A	9/1993	Miller et al.
4,939,310 A	7/1990	Wade	5,243,114 A	9/1993	Johnson et al.
4,939,311 A	7/1990	Washecheck et al.	5,245,109 A	9/1993	Kaminsky et al.
4,945,175 A	7/1990	Hobbs et al.	5,254,772 A	10/1993	Dukat et al.
4,950,811 A	8/1990	Doussain et al.	5,254,790 A	10/1993	Thomas et al.
4,950,822 A	8/1990	Dileo et al.	5,264,635 A	11/1993	Le et al.
4,956,521 A	9/1990	Volles	5,268,518 A	12/1993	West et al.
4,962,252 A	10/1990	Wade	5,276,226 A	1/1994	Horvath et al.
4,973,776 A	11/1990	Allenger et al.	5,276,240 A	1/1994	Timmons et al.
4,973,786 A	11/1990	Karra	5,276,242 A	1/1994	Wu
4,982,024 A	1/1991	Lin et al.	5,284,990 A	2/1994	Peterson et al.
4,982,041 A	1/1991	Campbell	5,300,126 A	4/1994	Brown et al.
4,988,660 A	1/1991	Campbell	5,306,855 A	4/1994	Periana et al.
4,990,696 A	2/1991	Stauffer	5,316,995 A	5/1994	Kaminsky et al.
4,990,711 A	2/1991	Chen et al.	5,319,132 A	6/1994	Ozawa et al.
5,001,293 A	3/1991	Nubel et al.	5,334,777 A	8/1994	Miller et al.
5,004,847 A	4/1991	Beaver et al.	5,345,021 A	9/1994	Casci et al.
5,013,424 A	5/1991	James, Jr. et al.	5,354,916 A	10/1994	Horvath et al.
5,013,793 A	5/1991	Wang et al.	5,354,931 A	10/1994	Jan et al.
5,019,652 A	5/1991	Taylor et al.	5,366,949 A	11/1994	Schubert
5,026,934 A	6/1991	Bains et al.	5,371,313 A	12/1994	Ostrowicki
5,026,937 A	6/1991	Bricker	5,382,704 A	1/1995	Krespan et al.
5,026,944 A	6/1991	Allenger et al.	5,382,743 A	1/1995	Beech, Jr. et al.
5,034,566 A	7/1991	Ishino et al.	5,382,744 A	1/1995	Abbott et al.
5,043,502 A	8/1991	Martindale et al.	5,385,718 A	1/1995	Casci et al.
5,055,235 A	10/1991	Brackenridge et al.	5,395,981 A	3/1995	Marker
5,055,633 A	10/1991	Volles	5,399,258 A	3/1995	Fletcher et al.
5,055,634 A	10/1991	Volles	5,401,890 A	3/1995	Parks
5,059,744 A	10/1991	Harandi et al.	5,401,894 A	3/1995	Brasier et al.
5,068,478 A	11/1991	Miller et al.	5,406,017 A	4/1995	Withers, Jr.
5,071,449 A	12/1991	Sircar	5,414,173 A	5/1995	Garces et al.
5,071,815 A	12/1991	Wallace	5,430,210 A	7/1995	Grasselli et al.
5,073,656 A	12/1991	Chafin et al.	5,430,214 A	7/1995	Smith et al.
5,073,657 A	12/1991	Warren	5,430,219 A	7/1995	Sanfilippo et al.

# US 7,838,708 B2

Page 4

---

5,436,378 A	7/1995	Masini et al.	5,983,476 A	11/1999	Eshelman et al.
5,444,168 A	8/1995	Brown	5,986,158 A	11/1999	Van Broekhoven et al.
5,446,234 A	8/1995	Casci et al.	5,994,604 A	11/1999	Reagen et al.
5,453,557 A	9/1995	Harley et al.	5,998,679 A	12/1999	Miller
5,456,822 A	10/1995	Marcilly et al.	5,998,686 A	12/1999	Clem et al.
5,457,255 A	10/1995	Kumata et al.	6,002,059 A	12/1999	Hellring et al.
5,464,799 A	11/1995	Casci et al.	6,015,867 A	1/2000	Fushimi et al.
5,465,699 A	11/1995	Voigt	6,018,088 A	1/2000	Olah
5,470,377 A	11/1995	Whitlock	6,022,929 A	2/2000	Chen et al.
5,480,629 A	1/1996	Thompson et al.	6,034,288 A	3/2000	Scott et al.
5,486,627 A	1/1996	Quarderer et al.	6,056,804 A	5/2000	Keefer et al.
5,489,719 A	2/1996	Le et al.	6,068,679 A	5/2000	Zheng
5,489,727 A	2/1996	Randolph et al.	6,072,091 A	6/2000	Cosyns et al.
5,500,297 A	3/1996	Thompson et al.	6,087,294 A	7/2000	Klabunde et al.
5,510,525 A	4/1996	Sen et al.	6,090,312 A	7/2000	Ziaka et al.
5,523,503 A	6/1996	Funk et al.	6,096,932 A	8/2000	Subramanian
5,525,230 A	6/1996	Wrigley et al.	6,096,933 A	8/2000	Cheung et al.
5,538,540 A	7/1996	Whitlock	6,103,215 A	8/2000	Zones et al.
5,563,313 A	10/1996	Chung et al.	6,107,561 A	8/2000	Thompson
5,565,092 A	10/1996	Pannell et al.	6,117,371 A	9/2000	Mack
5,565,616 A	10/1996	Li et al.	6,124,514 A	9/2000	Emmrich et al.
5,571,762 A	11/1996	Clerici et al.	6,127,588 A	10/2000	Kimble et al.
5,571,885 A	11/1996	Chung et al.	6,130,260 A	10/2000	Hall et al.
5,599,381 A	2/1997	Whitlock	6,143,939 A	11/2000	Farcasiu et al.
5,600,043 A	2/1997	Johnston et al.	6,169,218 B1	1/2001	Hearn et al.
5,600,045 A	2/1997	Van Der Aalst et al.	6,180,841 B1	1/2001	Fatutto et al.
5,609,654 A	3/1997	Le et al.	6,187,871 B1	2/2001	Thompson et al.
5,633,419 A	5/1997	Spencer et al.	6,187,983 B1	2/2001	Sun
5,639,930 A	6/1997	Penick	6,203,712 B1	3/2001	Bronner et al.
5,653,956 A	8/1997	Zones	6,207,864 B1	3/2001	Henningsen et al.
5,656,149 A	8/1997	Zones et al.	6,225,517 B1	5/2001	Nascimento et al.
5,661,097 A	8/1997	Spencer et al.	6,248,218 B1	6/2001	Linkous et al.
5,663,465 A	9/1997	Clegg et al.	6,265,505 B1	7/2001	McConville et al.
5,663,474 A	9/1997	Pham et al.	6,281,405 B1	8/2001	Davis et al.
5,675,046 A	10/1997	Ohno et al.	6,320,085 B1	11/2001	Arvai et al.
5,675,052 A	10/1997	Menon et al.	6,337,063 B1	1/2002	Rouleau et al.
5,679,134 A	10/1997	Brugeron et al.	6,342,200 B1	1/2002	Rouleau et al.
5,679,879 A	10/1997	Mercier et al.	6,368,490 B1	4/2002	Gestermann
5,684,213 A	11/1997	Nemphos et al.	6,369,283 B1	4/2002	Guram et al.
5,693,191 A	12/1997	Pividal et al.	6,372,949 B1	4/2002	Brown et al.
5,695,890 A	12/1997	Thompson et al.	6,376,731 B1	4/2002	Evans et al.
5,698,747 A	12/1997	Godwin et al.	6,380,328 B1	4/2002	McConville et al.
5,705,712 A	1/1998	Frey et al.	6,380,423 B2	4/2002	Banning et al.
5,705,728 A	1/1998	Viswanathan et al.	6,380,444 B1	4/2002	Bjerrum et al.
5,705,729 A	1/1998	Huang	6,395,945 B1	5/2002	Randolph
5,708,246 A	1/1998	Camaioni et al.	6,403,840 B1	6/2002	Zhou et al.
5,720,858 A	2/1998	Noceti et al.	6,406,523 B1	6/2002	Connor et al.
5,728,897 A	3/1998	Buyssch et al.	6,423,211 B1	7/2002	Randolph et al.
5,728,905 A	3/1998	Clegg et al.	6,426,441 B1	7/2002	Randolph et al.
5,734,073 A	3/1998	Chambers et al.	6,426,442 B1	7/2002	Ichikawa et al.
5,741,949 A	4/1998	Mack	6,452,058 B1	9/2002	Schweizer et al.
5,744,669 A	4/1998	Kalnes et al.	6,455,650 B1	9/2002	Lipian et al.
5,750,801 A	5/1998	Buyssch et al.	6,462,243 B1	10/2002	Zhou et al.
5,770,175 A	6/1998	Zones	6,465,696 B1	10/2002	Zhou et al.
5,776,871 A	7/1998	Cothran et al.	6,465,699 B1	10/2002	Grosso
5,780,703 A	7/1998	Chang et al.	6,472,345 B2	10/2002	Hintermann et al.
5,798,314 A	8/1998	Spencer et al.	6,472,572 B1	10/2002	Zhou et al.
5,814,715 A	9/1998	Chen et al.	6,475,463 B1	11/2002	Elomari et al.
5,817,904 A	10/1998	Vic et al.	6,475,464 B1	11/2002	Rouleau et al.
5,821,394 A	10/1998	Schoebrechts et al.	6,479,705 B2	11/2002	Murata et al.
5,847,224 A	12/1998	Koga et al.	6,482,997 B2	11/2002	Petit-Clair et al.
5,849,978 A	12/1998	Benazzi et al.	6,486,368 B1	11/2002	Zhou et al.
5,866,735 A	2/1999	Cheung et al.	6,495,484 B1	12/2002	Holtcamp
5,895,831 A	4/1999	Brasier et al.	6,509,485 B2	1/2003	Mul et al.
5,898,086 A	4/1999	Harris	6,511,526 B2	1/2003	Jagger et al.
5,905,169 A	5/1999	Jacobson	6,514,319 B2	2/2003	Keefer et al.
5,906,892 A	5/1999	Thompson et al.	6,518,474 B1	2/2003	Sanderson et al.
5,908,963 A	6/1999	Voss et al.	6,518,476 B1	2/2003	Culp et al.
5,952,538 A	9/1999	Vaughn et al.	6,525,228 B2	2/2003	Chauvin et al.
5,959,170 A	9/1999	Withers	6,525,230 B2	2/2003	Grosso
5,968,236 A	10/1999	Bassine	6,528,693 B1	3/2003	Gandy et al.
5,969,195 A	10/1999	Stabel et al.	6,538,162 B2	3/2003	Chang et al.
5,977,402 A	11/1999	Sekiguchi et al.	6,540,905 B1	4/2003	Elomari

# US 7,838,708 B2

Page 5

---

6,545,191	B1	4/2003	Stauffer	7,091,270	B2	8/2006	Zilberman et al.
6,547,958	B1	4/2003	Elomari	7,091,387	B2	8/2006	Fong et al.
6,548,040	B1	4/2003	Rouleau et al.	7,091,391	B2	8/2006	Stauffer
6,552,241	B1	4/2003	Randolph et al.	7,094,936	B1	8/2006	Owens et al.
6,566,572	B2	5/2003	Okamoto et al.	7,098,371	B2	8/2006	Mack et al.
6,572,829	B2	6/2003	Linkous et al.	7,105,710	B2	9/2006	Boons et al.
6,585,953	B2	7/2003	Roberts et al.	7,138,534	B2	11/2006	Forlin et al.
6,616,830	B2	9/2003	Elomari	7,141,708	B2	11/2006	Marsella et al.
6,620,757	B2	9/2003	McConville et al.	7,145,045	B2	12/2006	Harmsen et al.
6,632,971	B2	10/2003	Brown et al.	7,148,356	B2	12/2006	Smith, III et al.
6,635,793	B2	10/2003	Mul et al.	7,148,390	B2	12/2006	Zhou et al.
6,641,644	B2	11/2003	Jagger et al.	7,151,199	B2	12/2006	Martens et al.
6,646,102	B2	11/2003	Boriack et al.	7,161,050	B2	1/2007	Sherman et al.
6,669,846	B2	12/2003	Perriello	7,169,730	B2	1/2007	Ma et al.
6,672,572	B2	1/2004	Werlen	7,176,340	B2	2/2007	Van Broekhoven et al.
6,679,986	B1	1/2004	Da Silva et al.	7,176,342	B2	2/2007	Bellussi et al.
6,680,415	B1	1/2004	Gulotty, Jr. et al.	7,182,871	B2	2/2007	Perriello
6,692,626	B2	2/2004	Keefer et al.	7,193,093	B2	3/2007	Murray et al.
6,692,723	B2	2/2004	Rouleau et al.	7,196,239	B2	3/2007	Van Egmond et al.
6,710,213	B2	3/2004	Aoki et al.	7,199,083	B2	4/2007	Zevallos
6,713,087	B2	3/2004	Tracy et al.	7,199,255	B2	4/2007	Murray et al.
6,713,655	B2	3/2004	Yilmaz et al.	7,208,641	B2	4/2007	Nagasaki et al.
RE38,493	E	4/2004	Keefer et al.	7,214,750	B2	5/2007	McDonald et al.
6,723,808	B2	4/2004	Holtcamp	7,220,391	B1	5/2007	Huang et al.
6,727,400	B2	4/2004	Messier et al.	7,226,569	B2	6/2007	Elomari
6,740,146	B2	5/2004	Simonds	7,226,576	B2	6/2007	Elomari
6,753,390	B2	6/2004	Ehrman et al.	7,230,150	B2	6/2007	Grosso et al.
6,765,120	B2	7/2004	Weber et al.	7,230,151	B2	6/2007	Martens et al.
6,797,845	B1	9/2004	Hickman et al.	7,232,872	B2	6/2007	Shaffer et al.
6,797,851	B2	9/2004	Martens et al.	7,238,846	B2	7/2007	Janssen et al.
6,821,924	B2	11/2004	Gulotty, Jr. et al.	7,244,795	B2	7/2007	Agapiou et al.
6,822,123	B2	11/2004	Stauffer	7,244,867	B2	7/2007	Waycuilis
6,822,125	B2	11/2004	Lee et al.	7,250,107	B2	7/2007	Benazzi et al.
6,825,307	B2	11/2004	Goodall	7,250,542	B2	7/2007	Smith, Jr. et al.
6,825,383	B1	11/2004	Dewkar et al.	7,252,920	B2	8/2007	Kurokawa et al.
6,831,032	B2	12/2004	Spaether	7,253,327	B2	8/2007	Janssens et al.
6,838,576	B1	1/2005	Wicki et al.	7,253,328	B2	8/2007	Stauffer
6,841,063	B2	1/2005	Elomari	7,265,193	B2	9/2007	Weng et al.
6,852,896	B2	2/2005	Stauffer	7,267,758	B2	9/2007	Benazzi et al.
6,866,950	B2	3/2005	Connor et al.	7,268,263	B1	9/2007	Frey et al.
6,869,903	B2	3/2005	Matsunaga	7,271,303	B1	9/2007	Sechrist et al.
6,875,339	B2	4/2005	Rangarajan et al.	7,273,957	B2	9/2007	Bakshi et al.
6,878,853	B2	4/2005	Tanaka et al.	7,282,603	B2	10/2007	Richards
6,888,013	B2	5/2005	Paparatto et al.	7,285,698	B2	10/2007	Liu et al.
6,900,363	B2	5/2005	Harth et al.	7,304,193	B1	12/2007	Frey et al.
6,902,602	B2	6/2005	Keefer et al.	7,342,144	B2	3/2008	Kaizik et al.
6,903,171	B2	6/2005	Rhodes et al.	7,348,295	B2	3/2008	Zones et al.
6,909,024	B1	6/2005	Jones et al.	7,348,464	B2	3/2008	Waycuilis
6,921,597	B2	7/2005	Keefer et al.	7,357,904	B2	4/2008	Zones et al.
6,933,417	B1	8/2005	Henley et al.	7,361,794	B2	4/2008	Grosso
6,946,566	B2	9/2005	Yaegashi et al.	7,390,395	B2	6/2008	Elomari
6,953,868	B2	10/2005	Boaen et al.	2002/0102672	A1	8/2002	Mizrahi
6,953,873	B2	10/2005	Cortright et al.	2002/0198416	A1	12/2002	Zhou et al.
6,956,140	B2	10/2005	Ehrenfeld	2003/0004380	A1	1/2003	Grumann
6,958,306	B2	10/2005	Holtcamp	2003/0065239	A1	4/2003	Zhu
6,984,763	B2	1/2006	Schweizer et al.	2003/0069452	A1	4/2003	Sherman et al.
7,001,872	B2	2/2006	Pyecroft et al.	2003/0078456	A1	4/2003	Yilmaz et al.
7,002,050	B2	2/2006	Santiago Fernandez et al.	2003/0120121	A1	6/2003	Sherman et al.
7,011,811	B2	3/2006	Elomari	2003/0125589	A1	7/2003	Grosso
7,019,182	B2	3/2006	Grosso	2003/0166973	A1	9/2003	Zhou et al.
7,026,145	B2	4/2006	Mizrahi et al.	2004/0006246	A1	1/2004	Sherman et al.
7,026,519	B2	4/2006	Santiago Fernandez et al.	2004/0062705	A1	4/2004	Leduc
7,037,358	B2	5/2006	Babicki et al.	2004/0152929	A1	8/2004	Clarke
7,045,670	B2	5/2006	Johnson et al.	2004/0158107	A1	8/2004	Aoki
7,049,388	B2	5/2006	Boriack et al.	2004/0158108	A1	8/2004	Snoble
7,053,252	B2	5/2006	Boussand et al.	2004/0187684	A1	9/2004	Elomari
7,057,081	B2	6/2006	Allison et al.	2005/0047927	A1	3/2005	Lee et al.
7,060,865	B2	6/2006	Ding et al.	2005/0148805	A1	7/2005	Jones
7,064,238	B2	6/2006	Waycuilis	2005/0171393	A1	8/2005	Lorkovic
7,064,240	B2	6/2006	Ohno et al.	2005/0192468	A1	9/2005	Sherman et al.
7,067,448	B1	6/2006	Weitkamp et al.	2005/0215837	A1	9/2005	Hoffpauir
7,083,714	B2	8/2006	Elomari	2005/0234276	A1	10/2005	Waycuilis
7,084,308	B1	8/2006	Stauffer	2005/0245772	A1	11/2005	Fong

2005/0245777	A1	11/2005	Fong	GB	156122	3/1922
2005/0267224	A1	12/2005	Herling	GB	294100	6/1929
2006/0025617	A1	2/2006	Begley	GB	363009	12/1931
2006/0100469	A1	5/2006	Waycuilis	GB	402928	12/1933
2006/0135823	A1	6/2006	Jun	GB	474922 A	11/1937
2006/0138025	A1	6/2006	Zones	GB	536491	5/1941
2006/0138026	A1	6/2006	Chen	GB	553950	6/1943
2006/0149116	A1	7/2006	Slaugh	GB	586483	3/1947
2006/0229228	A1	10/2006	Komon et al.	GB	775590	5/1957
2006/0229475	A1	10/2006	Weiss et al.	GB	793214	4/1958
2006/0270863	A1	11/2006	Reiling	GB	796048	6/1958
2006/0288690	A1	12/2006	Elomari	GB	796085	6/1958
2007/0004955	A1	1/2007	Kay	GB	883256	11/1961
2007/0078285	A1	4/2007	Dagle	GB	950975	3/1964
2007/0100189	A1	5/2007	Stauffer	GB	950976	3/1964
2007/0129584	A1	6/2007	Basset	GB	991303	5/1965
2007/0142680	A1	6/2007	Ayoub	GB	995960	6/1965
2007/0148067	A1	6/2007	Zones	GB	1015033	12/1965
2007/0148086	A1	6/2007	Zones	GB	1104294	2/1968
2007/0149778	A1	6/2007	Zones	GB	1133752	11/1968
2007/0149789	A1	6/2007	Zones	GB	1172002	11/1969
2007/0149819	A1	6/2007	Zones	GB	1212240	11/1970
2007/0149824	A1	6/2007	Zones	GB	1233299	5/1971
2007/0149837	A1	6/2007	Zones	GB	1253618	11/1971
2007/0197801	A1	8/2007	Bolk	GB	1263806 A	2/1972
2007/0197847	A1	8/2007	Liu	GB	1446803	8/1976
2007/0213545	A1	9/2007	Bolk	GB	1542112	3/1979
2007/0238905	A1	10/2007	Arredondo	GB	2095243 A	9/1982
2007/0238909	A1	10/2007	Gadewar et al.	GB	2095245 A	9/1982
2007/0251382	A1	11/2007	Gadewar	GB	2095249 A	9/1982
2007/0276168	A1	11/2007	Garel	GB	2116546 A	9/1982
2007/0284284	A1	12/2007	Zones	GB	2120249 A	11/1983
2008/0171898	A1	7/2008	Waycuilis	GB	2185754 A	7/1987
2008/0183022	A1	7/2008	Waycuilis	GB	2191214 A	12/1987
2008/0188697	A1	8/2008	Lorkovic	JP	2004-529189	9/2004
2008/0269534	A1	10/2008	Lorkovic	WO	83/00859	3/1983
2008/0314758	A1	12/2008	Grosso	WO	85/04863	11/1985
2009/0069606	A1	3/2009	Komon	WO	85/04867	11/1985
2009/0127163	A1	5/2009	Weiss	WO	90/08120	7/1990
2010/0096588	A1	4/2010	Gadewar	WO	90/08752	8/1990
2010/0099928	A1	4/2010	Gadewar	WO	91/18856	12/1991
2010/0099929	A1	4/2010	Gadewar	WO	92/03401	3/1992
2010/0099930	A1	4/2010	Stoimenov	WO	92/12946	8/1992
2010/0105972	A1	4/2010	Lorkovic	WO	93/16798	9/1993
				WO	96/22263	7/1996
				WO	97/44302	11/1997
				WO	98/12165	3/1998

## FOREIGN PATENT DOCUMENTS

EP	0021497	1/1981	WO	99/07443	2/1999
EP	0164798 A1	12/1985	WO	00/07718 A1	2/2000
EP	0418971 A1	3/1991	WO	00/09261 A1	2/2000
EP	0418974 A1	3/1991	WO	01/14300 A1	3/2001
EP	0418975 A1	3/1991	WO	01/38275 A1	5/2001
EP	0510238 A1	10/1992	WO	01/44149 A1	6/2001
EP	0526908 A2	2/1993	WO	02/094749 A1	11/2002
EP	0346612 B1	8/1993	WO	02/094750 A1	11/2002
EP	0560546 A1	9/1993	WO	02/094751 A2	11/2002
EP	0976705 A1	7/1998	WO	02/094752 A1	11/2002
EP	1186591 A2	3/2002	WO	03/000635 A1	1/2003
EP	1253126 A1	10/2002	WO	03/002251 A2	1/2003
EP	1312411 A2	5/2003	WO	03/018524 A1	3/2003
EP	1395536	3/2004	WO	03/020676 A1	3/2003
EP	1404636	4/2004	WO	03/022827 A1	3/2003
EP	1235769 B1	5/2004	WO	03/043575 A2	5/2003
EP	1435349 A2	7/2004	WO	03/051813 A1	6/2003
EP	1440939 A1	7/2004	WO	03/062143 A1	7/2003
EP	1474371	11/2004	WO	03/062172 A2	7/2003
EP	1235772 B1	1/2005	WO	03/078366 A1	9/2003
EP	1661620 A1	5/2006	WO	2004/018093 A2	3/2004
EP	1760057 A1	3/2007	WO	2004/067487 A2	8/2004
EP	1689728 B1	4/2007	WO	2005/014168 A1	2/2005
EP	1808227 A1	7/2007	WO	2005/019143 A1	3/2005
EP	1837320 A1	9/2007	WO	2005/021468 A1	3/2005
GB	5125	2/1912	WO	2005/035121 A2	4/2005

WO	2005/037758	A1	4/2005	Abstract of WO0105737, Method for preparing a carboxylic acid, Publication date: Jan. 25, 2001, Inventor: Pascal et al., esp@cenet database—worldwide.
WO	2005/054120	A2	6/2005	Abstract of WO0105738, Method for preparing a carboxylic acid, Publication date: Jan. 25, 2001, Inventor: Pascal et al., esp@cenet database—worldwide.
WO	2005/056525	A2	6/2005	Abstract of WO2004092099, Method for producing cyclic enols, Publication date: Oct. 28, 2004, Inventor: Marko et al., esp@cenet database—worldwide.
WO	2005/058782	A1	6/2005	Abstract of WO2006063852, Electroluminescent polymers and use thereof, Publication date: Jun. 22, 2006, Inventor: Arne et al., esp@cenet database—worldwide.
WO	2005/090272	A1	9/2005	Abstract of WO2006136135, Method for decarboxylating C-C cross-linking of carboxylic acids with carbon electrophiles, Publication date: Dec. 28, 2006, Inventor: Goossen Lukas et al., esp@cenet database—worldwide.
WO	2005/095310	A2	10/2005	Abstract of WO2007028761, Method for chlorinating alcohols, Publication date: Mar. 15, 2007, Inventor: Rohde et al., esp@cenet database—worldwide.
WO	2005/105709	A1	11/2005	Abstract of WO2007128842, Catalytic transalkylation of dialkyl benzenes, Publication date: Nov. 15, 2007, Inventor: Goncalvesalmeida et al., esp@cenet database—worldwide.
WO	2005/105715	A1	11/2005	Abstract of WO2007137566, Method for catalytic conversion of organic oxygenated compounds from biomaterials, Publication date: Dec. 6, 2007, Inventor: Reschetilowski, esp@cenet database—worldwide.
WO	2005/110953	A1	11/2005	Abstract of WO9721656, Method for making fluoroalkanols, Publication date: Jun. 19, 1997, Inventor: Gillet, esp@cenet database—worldwide.
WO	2005/113437	A1	12/2005	Abstract of WO9950213, Method for producing dialkyl ethers, Publication date: Oct. 7, 1999, Inventor: Falkowski Juergen et al., esp@cenet database—worldwide.
WO	2005/113440	A1	12/2005	Adachi, et al.; Synthesis of Sialyl Lewis X Ganglioside Analogs Containing a Variable Length Spacer Between the Sugar and Lipophilic Moieties; <i>J. Carbohydrate Chem.</i> , vol. 17, No. 4-5, (1998), pp. 595-607, XP009081720.
WO	2006/007093	A1	1/2006	Abstract of EP0039471, Process for the preparation of 2-chloro-1,1,1,2,3,3,3-heptafluoropropane, Publication date: Nov. 11, 1981, Inventor: Von Halasz, esp@cenet database—worldwide.
WO	2006/015824	A1	2/2006	Abstract of EP0101337, Process for the production of methylene chloride, Publication date: Feb. 22, 1984, Inventor: Olah et al., esp@cenet database—worldwide.
WO	2006/019399	A2	2/2006	Abstract of EP0407989, Method for the production of 1,1,1-trifluoro-2,2-dichloroethane by photochlorination, Publication date: Jan. 16, 1991, Inventor: Cremer et al., esp@cenet database—worldwide.
WO	2006/020234	A1	2/2006	Abstract of EP0442258, Process for the preparation of a polyunsaturated olefin, Publication date: Aug. 21, 1991, Inventor: Gaudin et al., esp@cenet database—worldwide.
WO	2006/036293	A1	4/2006	Abstract of EP0465294, Process for the preparation of unsaturated bromides, Publication date: Jan. 8, 1992, Inventor: Decaudin et al., esp@cenet database—worldwide.
WO	2006/039213	A1	4/2006	Abstract of EP0549387, Synthesis of n-perfluoroctylbromide, Publication date: Jun. 30, 1993, Inventor: Drivon et al., esp@cenet database—worldwide.
WO	2006/039354	A2	4/2006	Abstract of EP0850906, Process and apparatus for the etherification of olefinic hydrocarbon feedstocks, Publication date: Jul. 1, 1998, Inventor: Masson, esp@cenet database—worldwide.
WO	2006/043075	A1	4/2006	Abstract of EP0858987, Process for the conversion of lighter alkanes to higher hydrocarbons, Publication date: Aug. 19, 1998, Inventor: Amariglio et al., esp@cenet database—worldwide.
WO	2006/053345	A1	5/2006	Bakker, et al.; An Exploratory Study of the Addition Reaction of Ethylenegcol, 2-Chloroethanol and 1, 3-Dichloro-2-Propanol to 1-Dodecene; <i>J. Am. Oil Chem. Soc.</i> , vol. 44, No. 9 (1967), pp. 517-521; XP009081570.
WO	2006-067155	A2	6/2006	Abstract of EP0235110, Process for the stabilization of silicalite catalysts, Publication date: Sep. 2, 1987, Inventor: Debras et al., esp@cenet database—worldwide.
WO	2006/067188	A1	6/2006	Bouzide et al.; Highly Selective Silver (I) Oxide Mediated Monoprotection of Symmetric Diols; <i>Tetrahedron Letters</i> , Elsevier, vol. 38, No. 34 (1997), pp. 5945-5948; XP004094157.
WO	2006/067190	A1	6/2006	Combined International Search Report and Written Opinion Dated Apr. 17, 2007 for PCT/US06/13394, in the name of GRT, Inc.
WO	2006/067191	A1	6/2006	
WO	2006/067192	A1	6/2006	
WO	2006/067193	A1	6/2006	
WO	2006/069107	A2	6/2006	
WO	2006/071354	A1	7/2006	
WO	2006/076942	A1	7/2006	
WO	2006/083427	A1	8/2006	
WO	2006-100312	A2	9/2006	
WO	2006/104909	A2	10/2006	
WO	2006/104914	A1	10/2006	
WO	2006/111997	A1	10/2006	
WO	2006/113205	A2	10/2006	
WO	2006/118935	A2	11/2006	
WO	2007/001934	A2	1/2007	
WO	2007/017900	A2	2/2007	
WO	2007/044139	A1	4/2007	
WO	2007/046986	A2	4/2007	
WO	2007/050745	A1	5/2007	
WO	2007/071046	A1	6/2007	
WO	2007/079038	A2	7/2007	
WO	2007/091009	A2	8/2007	
WO	2007/094995	A2	8/2007	
WO	2007/107031	A1	9/2007	
WO	2007/111997	A2	10/2007	
WO	2007/114479	A1	10/2007	
WO	2007/125332	A1	11/2007	
WO	2007/130054	A1	11/2007	
WO	2007/130055	A1	11/2007	
WO	2007/1421295	A1	12/2007	
WO	2007/142745	A1	12/2007	
WO	2008/036562		3/2008	
WO	2008/036563		3/2008	
WO	2008/106319		9/2008	
WO	2008/157043		12/2008	
WO	2008/157044		12/2008	
WO	2008/157045		12/2008	
WO	2008/157046		12/2008	
WO	2008/157047		12/2008	

## OTHER PUBLICATIONS

Abstract of JP2007061594, Method for decomposing organohalogen compound and mobile decomposition system, Publication date: Mar. 15, 2007, Inventor: Koichi et al., esp@cenet database—worldwide.

Abstract of JP2007099729, Method for producing alpha-methylstyrene or cumene, Publication date: Apr. 19, 2007, Inventor: Toshio, esp@cenet database—worldwide.

Abstract of RO119778, Process for preparing perchloroethylene, Publication date: Mar. 30, 2005, Inventor: Horia et al., esp@cenet database—worldwide.

Abstract of WO0105737, Method for preparing a carboxylic acid, Publication date: Jan. 25, 2001, Inventor: Pascal et al., esp@cenet database—worldwide.

Abstract of WO0105738, Method for preparing a carboxylic acid, Publication date: Jan. 25, 2001, Inventor: Pascal et al., esp@cenet database—worldwide.

Abstract of WO2004092099, Method for producing cyclic enols, Publication date: Oct. 28, 2004, Inventor: Marko et al., esp@cenet database—worldwide.

Abstract of WO2006063852, Electroluminescent polymers and use thereof, Publication date: Jun. 22, 2006, Inventor: Arne et al., esp@cenet database—worldwide.

Abstract of WO2006136135, Method for decarboxylating C-C cross-linking of carboxylic acids with carbon electrophiles, Publication date: Dec. 28, 2006, Inventor: Goossen Lukas et al., esp@cenet database—worldwide.

Abstract of WO2007028761, Method for chlorinating alcohols, Publication date: Mar. 15, 2007, Inventor: Rohde et al., esp@cenet database—worldwide.

Abstract of WO2007128842, Catalytic transalkylation of dialkyl benzenes, Publication date: Nov. 15, 2007, Inventor: Goncalvesalmeida et al., esp@cenet database—worldwide.

Abstract of WO2007137566, Method for catalytic conversion of organic oxygenated compounds from biomaterials, Publication date: Dec. 6, 2007, Inventor: Reschetilowski, esp@cenet database—worldwide.

Abstract of WO9721656, Method for making fluoroalkanols, Publication date: Jun. 19, 1997, Inventor: Gillet, esp@cenet database—worldwide.

Abstract of WO9950213, Method for producing dialkyl ethers, Publication date: Oct. 7, 1999, Inventor: Falkowski Juergen et al., esp@cenet database—worldwide.

Adachi, et al.; Synthesis of Sialyl Lewis X Ganglioside Analogs Containing a Variable Length Spacer Between the Sugar and Lipophilic Moieties; *J. Carbohydrate Chem.*, vol. 17, No. 4-5, (1998), pp. 595-607, XP009081720.

Abstract of EP0039471, Process for the preparation of 2-chloro-1,1,1,2,3,3,3-heptafluoropropane, Publication date: Nov. 11, 1981, Inventor: Von Halasz, esp@cenet database—worldwide.

Abstract of EP0101337, Process for the production of methylene chloride, Publication date: Feb. 22, 1984, Inventor: Olah et al., esp@cenet database—worldwide.

Abstract of EP0407989, Method for the production of 1,1,1-trifluoro-2,2-dichloroethane by photochlorination, Publication date: Jan. 16, 1991, Inventor: Cremer et al., esp@cenet database—worldwide.

Abstract of EP0442258, Process for the preparation of a polyunsaturated olefin, Publication date: Aug. 21, 1991, Inventor: Gaudin et al., esp@cenet database—worldwide.

Abstract of EP0465294, Process for the preparation of unsaturated bromides, Publication date: Jan. 8, 1992, Inventor: Decaudin et al., esp@cenet database—worldwide.

Abstract of EP0549387, Synthesis of n-perfluoroctylbromide, Publication date: Jun. 30, 1993, Inventor: Drivon et al., esp@cenet database—worldwide.

Abstract of EP0850906, Process and apparatus for the etherification of olefinic hydrocarbon feedstocks, Publication date: Jul. 1, 1998, Inventor: Masson, esp@cenet database—worldwide.

Abstract of EP0858987, Process for the conversion of lighter alkanes to higher hydrocarbons, Publication date: Aug. 19, 1998, Inventor: Amariglio et al., esp@cenet database—worldwide.

Bakker, et al.; An Exploratory Study of the Addition Reaction of Ethylenegcol, 2-Chloroethanol and 1, 3-Dichloro-2-Propanol to 1-Dodecene; *J. Am. Oil Chem. Soc.*, vol. 44, No. 9 (1967), pp. 517-521; XP009081570.

Abstract of EP0235110, Process for the stabilization of silicalite catalysts, Publication date: Sep. 2, 1987, Inventor: Debras et al., esp@cenet database—worldwide.

Bouzide et al.; Highly Selective Silver (I) Oxide Mediated Monoprotection of Symmetric Diols; *Tetrahedron Letters*, Elsevier, vol. 38, No. 34 (1997), pp. 5945-5948; XP004094157.

Combined International Search Report and Written Opinion Dated Apr. 17, 2007 for PCT/US06/13394, in the name of GRT, Inc.

- Gibson; Phase-Transfer Synthesis of Monoalkyl Ethers of Oligoethylene Glycols; *Journal of Organic Chemistry*, vol. 45, No. 6 (1980) pp. 1095-1098; XP002427776.
- Klabunde, Kenneth J., et al., Changes in Texture and Catalytic Activity of Nanocrystalline MgO during Its Transformation to MgCl<sub>2</sub> in the Reaction with 1-Chlorobutane, *J. Phys. Chem. B* 2001, 105, 3937-3941. cited by other.
- Loiseau et al.; Multigram Synthesis of Well-Defined Extended Bifunctional Polyethylene Glycol (PEG) Chains; *J. of Organic Chem.*, vol. 69, No. 3 (2004), pp. 639-647; XP002345040.
- Mihai et al.; Application of Bronsted-type LFER in the study of phospholipase C mechanism; *J. Am. Chem. Soc.*, vol. 125, No. 11 (2003) pp. 3236-3242; XP002427799.
- Motupally et al., Recycling Chlorine from Hydrogen Chloride: A New and Economical Electrolytic Process, *The Electrochemical Society Interface*, Fall 1998.
- Nishikawa et al.; Ultrasonic Relaxations in Aqueous Solutions of Alcohols and the Balance Between Hydrophobicity and Hydrophilicity of the Solutes; *J. Phys. Chem.* vol. 97, No. 14 (1993), pp. 3539-3544; XP002427775.
- Prelog et al.; Chirale 2,2'-Polyoxaalkano-9,9'-Spirobifluorene; *Helvetica Chimica ACTA*, vol. 62, No. 7, (1979) pp. 2285-2302; XP002030901.
- Shimizu et al., Gas-Phase Electrolysis of Hydrobromic Acid Using PTFE-Bonded Carbon Electrode, *Int. J. Hydrogen Energy*, vol. 13, No. 6, pp. 345-349, 1988.
- Velzen et al., HBr Electrolysis in the Ispra Mark 13A Flue Gas Desulphurization Process: Electrolysis in a DEM Cell, *J. of Applied Electrochemistry*, vol. 20, pp. 60-68, 1990.
- Whitesides et al.; Nuclear Magnetic Resonance Spectroscopy. The Effect of Structure on Magnetic Nonequivalence Due to Molecular Asymmetry; *J. Am. Chem. Soc.*, vol. 86, No. 13 (1964), pp. 2628-2634; XP002427774.
- JLM Technology Ltd.; "The Miller GLS Technology for Conversion of Light Hydrocarbons to Alcohols"; New Science for the Benefit of Humanity; May 31, 2000; pp. 1-10.
- Jaumain, Denis and Su, Bao-Lian; "Direct Catalytic Conversion of Chloromethane to Higher Hydrocarbons Over Various Protonic and Cationic Zeolite Catalysts as Studied by in-situ FTIR and Catalytic Testing"; 2000; pp. 1607-1612; *Studies in Surface Science and Catalysis* 130; Elsevier Science B.V.
- Taylor, Charles E.; "Conversion of Substituted Methanes Over ZSM-Catalysts"; 2000; pp. 3633-3638; *Studies in Surface Science and Catalysis* 130; Elsevier Science B.V.
- ZSM-5 Catalyst; <http://chemelab.ucsd.edu/methanol/memos/ZSM-5.html>; Nov. 6, 2003; p. 1.
- Final Report; "Abstract"; <http://chemelab.ucsd.edu/methanol/memos/final.html>; May 9, 2004; pp. 1-7.
- Driscoll, Daniel J.; "Direct Methane Conversion"; Federal Energy Technology Center, U.S. Department of Energy; M970779; 2001; pp. 1-10.
- Olah et al.; "Selective Monohalogenation of Methane over Supported Acid or Platinum Metal Catalysts and Hydrolysis of Methyl Halides . . ."; *J. American Chemical Society* 1985, vol. 107, 0002-7863/85/1507-7097\$01.50/0; pp. 7097-7105.
- Murray et al.; "Conversion of Methyl Halides to Hydrocarbons on Basic Zeolites: A Discovery by in Situ NMR"; *J. American Chemical Society* 1993, vol. 115; pp. 4732-4741.
- Lorkovic et al.; "A Novel Integrated Process for the Functionalization of Methane and Ethane: Bromine as Mediator", *Catalysis Today* 98; 2004; pp. 317-322.
- Lorkovic et al.; "C1 Oxidative Coupling via Bromine Activation and Tandem Catalytic Condensation over CaO/Zeolite Composites II . . ."; *Catalysis Today* 98; 2004; pp. 589-594.
- Olah et al.; "Antimony Pentafluoride/Graphite Catalyzed Oxidative Conversion of Methyl Halides with Copper Oxides (or Copper/Oxygen) to Dimethyl Ether"; *J. Org. Chem.* 1990, 55; 1990 American Chemical Society; pp. 4289-4293.
- Taylor, Charles E. et al.; "Direct Conversion of Methane to Liquid Hydrocarbons Through Chlorocarbon Intermediates"; 1988 Elsevier Science Publishers B.V. Amsterdam, Netherlands; pp. 483-489.
- Chang, Clarence D. et al.; "The Conversion of Methanol and Other O-Compounds to Hydrocarbons over Zeolite Catalysts"; *Journal of Catalysis* 47; 1977; Academic Press, Inc.; pp. 249-259.
- Zhou, Xiao-Ping et al.; "An Integrated Process for Partial Oxidation of Alkanes"; *Chem. Commun.* 2003; The Royal Society of Chemistry 2003; pp. 2294-2295.
- Sun, Shouli et al.; "A General Integrated Process for Synthesizing Olefin Oxides"; *Chem. Commun.* 2004; The Royal Society of Chemistry 2004; pp. 2100-2101.
- Lorkovic, Ivan M. et al.; "C1 Oxidative Coupling via Bromine Activation and Tandem Catalytic Condensation and Neutralization over CaO/Zeolite Composites II . . ."; *Catalysis Today* 98; 2004; pp. 589-594.
- Yilmaz, Aysen et al.; "Bromine Mediated Partial Oxidation of Ethane over Nanostructured Zirconia Supported Metal Oxide/Bromide"; *Microporous and Mesoporous Materials*, 79; 2005; pp. 205-214.
- Taylor, Charles E.; "PETC's On-Site Natural Gas Conversion Efforts"; *Preprints of the Fuel Division, 208th National Meeting of the American Chemical Society*, 39 (4); 1994; pp. 1228-1232.
- Ione et al.; "Syntheses of Hydrocarbons from Compounds Containing One Carbon Atom Using Bifunctional Zeolite Catalysts"; *Solid Fuel Chemistry (Khimiya Tverdogo Topliva)*; 1982; pp. 29-43; vol. 16, No. 6; Allerton Press, Inc.
- Olah, George A. et al.; "Hydrocarbons Through Methane Derivatives"; *Hydrocarbon Chemistry*; 1995; pp. 89-90; John Wiley & Sons, Inc.
- Akhrem, Irena S. et al.; "Ionic Bromination of Ethane and Other Alkanes (Cycloalkanes) with Bromine Catalyzed by the Polyhalomethane-2AlBr<sub>3</sub> Aprotic Organic Superacids Under Mild Conditions"; *Tetrahedron Letters*, vol. 36, No. 51, 1995; pp. 9365-9368; Pergamon; Great Britain.
- Smirnov, Vladimir V. et al.; "Selective Bromination of Alkanes and Arylalkanes with CBr<sup>4</sup>"; *Mendeleev Commun.* 2000; pp. 175-176.
- Olah, George A.; "Electrophilic Methane Conversion"; *Acc. Chem. Res.* 1987, 20; pp. 422-428; American Chemical Society, Loker Hydrocarbon Research Institute and Dept. Of Chemistry; University of Southern California.
- Olah, George A. et al.; "Antimony Pentafluoride/Graphite Catalyzed Oxidative Carbonylation of Methyl Halides with Carbon Monoxide and Copper Oxides (or Copper/Oxygen) to Methyl Acetate"; *J. Org. Chem.* 1990, 55; pp. 4293-4297; Loker Hydrocarbon Research Institute and Dept. of Chemistry; University of Southern California.
- Bagno, Alessandro et al.; "Superacid-Catalyzed Carbonylation of Methane, Methyl Halides, Methyl Alcohol, and Dimethyl Ether to Methyl Acetate and Acetic Acid"; *J. Org. Chem.* 1990, 55; pp. 4284-4289; Loker Hydrocarbon Research Institute; University of Southern California.
- Olah, George A. et al.; "Onium Ylide Chemistry. 1. Bifunctional Acid-Base-Catalyzed Conversion of Heterosubstituted Methanes into Ethylene and Derived Hydrocarbons. The Onium Ylide Mechanism of the C1-C2 Conversion"; *J. Am. Chem. Soc.* 1984, 106; pp. 2143-2149.
- Mochida, Isao et al.; "The Catalytic Dehydrohalogenation of Haloethanes on Solid Acids and Bases"; *Bulletin of the Chemical Society of Japan*, vol. 44; 1971; pp. 3305-3310.
- Richards, Ryan et al.; "Nanocrystalline Ultra High Surface Area Magnesium Oxide as a Selective Base Catalyst"; *Scripta Materialia*, 44; 2001; pp. 1663-1666; Elsevier Science Ltd.
- Sun, Naijian et al.; "Nanocrystal Metal Oxide—Chlorine Adducts: Selective Catalysts for Chlorination of Alkanes"; *J. Am. Chem. Soc.* 1999, 121; pp. 5587-5588; American Chemical Society.
- Mishakov, Ilya V. et al.; "Nanocrystalline MgO as a Dehydrohalogenation Catalyst"; *Journal of Catalysis* 206; 2002; pp. 40-48; Elsevier Science, USA.
- Wagner, George W. et al.; "Reactions of VX, GD, and HD with Nanosize CaO: Autocatalytic Dehydrohalogenation of HD"; *J. Phys. Chem. B* 2000, 104; pp. 5118-5123; 2000 American Chemical Society.
- Fenelonov, Vladimir B. et al.; "Changes in Texture and Catalytic Activity of Nanocrystalline MgO during Its Transformation to MgCl<sub>2</sub> in the Reaction with 1-Chlorobutane"; *J. Phys. Chem. B* 2001, 105; pp. 3937-3941; 2001 American Chemical Society.

- <http://webbook.nist.gov/>; "Welcome to the NIST Chemistry Web-Book"; 2005; U.S. Secretary of Commerce on Behalf of the United States of America.
- Claude, Marion C. et al.; "Monomethyl-Branching of Long n-Alkanes in the Range from Decane to Tetraicosane on Pt/H-ZSM-22 Bifunctional Catalyst"; Journal of Catalysis 190; 2000; pp. 39-48.
- Thomas, J. M. et al.; "Synthesis and Characterization of a Catalytically Active Nickel-Silicoaluminophosphate Catalyst for the Conversion of Methanol to Ethene"; Chem. Mater.; 1991, 3; pp. 667-672; 1991 American Chemical Society.
- Thomas, John Meurig et al.; "Catalytically Active Centres in Porous Oxides: Design and Performance of Highly Selective New Catalysts"; Chem. Commun.; 2001; pp. 675-687.
- Lorkovic, Ivan et al.; "C1 Coupling via Bromine Activation and Tandem Catalytic Condensation and Neutralization over CaO/Zeolite Composites"; Chem. Commun., 2004; pp. 566-567.
- Tamura, Masuhiko et al.; "The Reactions of Grignard Reagents with Transition Metal Halides: Coupling, Disproportionation, and Exchange with Olefins"; Bulletin of the Chemical Society of Japan, vol. 44.; Nov. 1971; pp. 3063-3073.
- Weissmer, Klaus et al.; "Industrial Organic Chemistry"; 3rd Edition 1997. pp. 160-162, and 208.
- Abstract of BE812868, Aromatic hydrocarbons prodn. from chlorinated hydrocarbons, Publication date: Sep. 27, 1974, esp@cenet database—worldwide.
- Abstract of BE814900, Volatile aromatic cpds. prodn., Publication date: Sep. 2, 1974, esp@cenet database—worldwide.
- Abstract of CN1199039, Pentanol and its production process, Publication date: Nov. 18, 1998, Inventor: Kailun, esp@cenet database—worldwide.
- Abstract of CN1210847, Process for producing low carbon alcohol by directly hydrating low carbon olefines, Publication date: Mar. 17, 1999, Inventor: Zhenguo et al., esp@cenet database—worldwide.
- Abstract of CN1321728, Method for preparing aromatic hydrocarbon and hydrogen gas by using low-pressure gas, Publication date: Nov. 11, 2001, Inventor: Jie et al., esp@cenet database—worldwide.
- Abstract of CN1451721, Process for non-catalytic combustion deoxidizing coal mine gas for producing methanol, Publication date: Oct. 29, 2003, Inventor: Pengwan et al., esp@cenet database—worldwide.
- Abstract of CN1623969, Method for preparing 1, 4-benzene dimethanol, Publication date: Jun. 8, 2005, Inventor: Jiarong et al., esp@cenet database—worldwide.
- Abstract of CN1657592, Method for converting oil to multiple energy fuel product, Publication date: Aug. 24, 2005, Inventor: Li, esp@cenet database—worldwide.
- Abstract of CN1687316, Method for producing biologic diesel oil from rosin, Publication date: Oct. 26, 2005, Inventor: Jianchun et al., esp@cenet database—worldwide.
- Abstract of CN1696248, Method for synthesizing biologic diesel oil based on ion liquid, Publication date: Nov. 16, 2005, Inventor: Sun, esp@cenet database—worldwide.
- Abstract of CN1699516, Process for preparing bio-diesel-oil by using microalgae fat, Publication date: Nov. 23, 2005, Inventor: Miao, esp@cenet database—worldwide.
- Abstract of CN1704392, Process for producing alkylbenzene, Publication date: Dec. 7, 2005, Inventor: Gao, esp@cenet database—worldwide.
- Abstract of CN1724612, Biological diesel oil catalyst and method of synthesizing biological diesel oil using sai catalyst, Publication date: Jan. 25, 2006, Inventor: Gu, esp@cenet database—worldwide.
- Abstract of CN1986737, Process of producing biodiesel oil with catering waste oil, Publication date: Jun. 27, 2007, Inventor: Chen, esp@cenet database—worldwide.
- Abstract of CN100999680, Esterification reaction tech. of preparing biodiesel by waste oil, Publication date: Jul. 18, 2007, Inventor: Weiming, esp@cenet database—worldwide.
- Abstract of CN101016229, Refining method for bromomeoamyl alcohol, Publication date: Aug. 15, 2007, Inventor: Tian, esp@cenet database—worldwide.
- Abstract of DE3209964, Process for the preparation of chlorinated hydrocarbons, Publication date: Nov. 11, 1982, Inventor: Pyke et al., esp@cenet database—worldwide.
- Abstract of DE3210196, Process for the preparation of a monochlorinated olefin, Publication date: Jan. 5, 1983, Inventor: Pyke et al., esp@cenet database—worldwide.
- Abstract of DE3226028, Process for the preparation of monochlorinated olefin, Publication date: Feb. 3, 1983, Inventor: Pyke et al., esp@cenet database—worldwide.
- Abstract of DE3334225, Process for the preparation of 1, 2-dichloroethane, Publication date: Apr. 4, 1985, Inventor: Hebgen et al., esp@cenet database—worldwide.
- Abstract of DE4232056, 2,5-Di:methyl-hexane-2,5-di:ol continuous prodn. from tert. butanol—by oxidative dimerisation in two phase system with vigorous stirring, using aq. phase with specified density to facilitate phase sepn., Publication date: Mar. 31, 1994, Inventor: Gnann et al., esp@cenet database—worldwide.
- Abstract of DE4434823, Continuous prodn. of hydroxy-benzyl alkyl ether, Publication date: Apr. 4, 1996, Inventor: Stein et al., esp@cenet database—worldwide.
- Abstract of FR2692259, Aromatisation of 2-4C hydrocarbons—using a fixed-mobile-catalytic bed process, Publication date: Dec. 17, 1993, Inventor: Alario et al., esp@cenet database—worldwide.
- Abstract of FR2880019, Manufacturing 1, 2-dichloroethane, comprises cracking core hydrocarbonated source, separating into fractions, sending into chlorination reaction chamber and oxychlorination reaction chamber and separating from chambers, Publication date: Jun. 30, 2006, Inventor: Strebelle et al., esp@cenet database—worldwide.
- Abstract of FR2883870, Formation of 1, 2-dichloroethane useful in manufacture of vinyl chloride involves subjecting mixture of cracking products obtained by cracking of hydrocarbon source, to a succession of aqueous quenching, alkaline washing, and oxidation steps, Publication date: Oct. 6, 2006, Inventor: Balthasar et al., esp@cenet database—worldwide.
- Abstract of FR2883871, Preparing 1, 2-dichloroethane comprises cracking hydrocarbon to form mixture, sending mixture into storage reservoir, supplying mixture into chlorination and/or oxychlorination reactor, and separating 1, 2-dichloroethane from reactor, Publication date: Oct. 6, 2006, Inventor: Balthasar et al., esp@cenet database—worldwide.
- Abstract of IT1255246, Process for the preparation of dinitrodiphenylmethanes, Publication date: Oct. 20, 1995, Applicant: Enichem Spa et al., esp@cenet database—worldwide.
- Abstract of IT1255358, Process for the synthesis of 1, 4-butanediol, Publication date: Oct. 31, 1995, Inventor: Marco, esp@cenet database—worldwide.
- Abstract of JP2142740, Production of fluoroalcohol, Publication date: May 31, 1990, Inventor: Tsutomu et al., esp@cenet database—worldwide.
- Abstract of JP2144150, Chemical process and catalyst used therefore, Publication date: Jun. 1, 1990, Inventor: Deidamus et al., esp@cenet database—worldwide.
- Abstract of JP4305542, Production of halogenated hydrocarbon compounds, Publication date: Oct. 28, 1992, Inventor: Shinsuke et al., esp@cenet database—worldwide.
- Abstract of JP6172225, Method for fluorinating halogenated hydrocarbon, Publication date: Jun. 21, 1994, Inventor: Takashi et al., esp@cenet database—worldwide.
- Abstract of JP6206834, Production of tetrachloroethanes, Publication date: Jul. 26, 1994, Inventor: Toshiro et al., esp@cenet database—worldwide.
- Abstract of JP8266888, Method for decomposing aromatic halogen compound, Publication date: Oct. 15, 1996, Inventor: Yuuji et al., esp@cenet database—worldwide.
- Abstract of JP2001031605, Production of 3-hydroxy-1-cycloalkene, Publication date: Feb. 6, 2001, Inventor: Hideo et al., esp@cenet database—worldwide.
- Abstract of JP2004075683, Method for producing optically active halogenohydroxypropyl compound and glycidyl compound, Publication date: Mar. 11, 2004, Inventor: Keisuke et al., esp@cenet database—worldwide.
- Abstract of JP2004189655, Method for fluorinating with microwave, Publication date: Jul. 8, 2004, Inventor: Masaharu et al., esp@cenet database—worldwide.

Abstract of JP2005075798, Method for Producing adamantyl ester compound, Publication date: Mar. 24, 2005, Inventor: Norihiro et al., esp@cenet database—worldwide.

Abstract of JP2005082563, Method for producing 1, 3-adamantanediol, Publication date: Mar. 31, 2005, Inventor: Norihiro et al., esp@cenet database—worldwide.

Abstract of JP2005145977, Process for catalytically oxidizing olefin and cycloolefin for the purpose of forming enol, olefin ketone, and epoxide, Publication date: Jun. 9, 2005, Inventor: Cancheng et al., esp@cenet database—worldwide.

Abstract of JP2005254092, Method of manufacturing alkynes, Publication date: Sep. 22, 2005, Inventor: Shirakawa Eiji, esp@cenet database—worldwide.

Abstract of JP2006151892, Preparation method of alcohol derivative, Publication date: Jun. 15, 2006, Inventor: Baba Akio et al., esp@cenet database—worldwide.

Abstract of JP2006152263, Organic-inorganic hybrid-type mesoporous material, method for producing the same, and solid catalyst, Publication date: Jun. 15, 2006, Inventor: Junko et al., esp@cenet database—worldwide.

Abstract of JP2006193473, Aryl polyadamantane derivative having carboxy or acid anhydride group and method for producing the same, Publication date: Jul. 27, 2006, Inventor: Yasuto et al., esp@cenet database—worldwide.

Abstract of JP2006231318, Phosphorus containing macromolecule immobilizing palladium catalyst and method for using the same, Publication date: Sep. 7, 2006, Inventor: Osamu et al., esp@cenet database—worldwide.

Abstract of JP2006263567, Optical resolution method of optical isomer and optical resolution device, Publication date: Oct. 5, 2006, Inventor: Yoshikazu et al., esp@oenet database—worldwide.

Abstract of JP2006265157, Method for catalytically activating silicated nucleating agent using phosphazene base, Publication date: Oct. 5, 2006, Inventor: Yoshinori et al., esp@cenet database—worldwide.

Abstract of JP2006306758, Method for producing biaryl compound, Publication date: Nov. 9, 2006, Inventor: Yuji et al., esp@cenet database—worldwide.

Abstract of JP2007001942, Production method of para-xylene, Publication date: Jan. 11, 2007, Inventor: Kazuyoshi, esp@cenet database—worldwide.

Abstract of JP2007015994, Method for synthesizing organic compound in ultra high rate under high temperature and high pressure water, and system of high temperature and high pressure reaction, Publication date: Jan. 25, 2007, Inventor: Hajime et al., esp@cenet database—worldwide.

U.S. Office Action from U.S. Appl. No. 10/826,885 dated Oct. 31, 2005.

U.S. Office Action from U.S. Appl. No. 10/826,885 dated Apr. 19, 2006.

U.S. Office Action from U.S. Appl. No. 10/826,885 dated Jul. 27, 2006.

U.S. Office Action from U.S. Appl. No. 10/826,885 dated Nov. 2, 2006.

U.S. Office Action from U.S. Appl. No. 10/826,885 dated Jan. 24, 2007.

U.S. Office Action from U.S. Appl. No. 11/101,886 dated Jan. 24, 2007.

U.S. Office Action from U.S. Appl. No. 11/254,438 dated Jan. 24, 2007.

U.S. Office Action from U.S. Appl. No. 11/254,438 dated Nov. 1, 2007.

U.S. Office Action from U.S. Appl. No. 10/893,418 dated Jan. 2, 2008.

U.S. Office Action from U.S. Appl. No. 10/893,418 dated Jun. 14, 2007.

U.S. Office Action from U.S. Appl. No. 11/091,130 dated Oct. 3, 2007.

U.S. Office Action from U.S. Appl. No. 10/365,346 dated Jun. 12, 2006.

U.S. Office Action from U.S. Appl. No. 11/103,326 dated Aug. 31, 2007.

U.S. Office Action from U.S. Appl. No. 11/103,326 dated Dec. 6, 2006.

U.S. Office Action from U.S. Appl. No. 11/098,997 dated Nov. 20, 2008.

U.S. Office Action from U.S. Appl. No. 12/215,326 dated Feb. 10, 2009.

U.S. Office Action from U.S. Appl. No. 10/430,240 dated Mar. 6, 2006.

U.S. Office Action from U.S. Appl. No. 10/369,148 dated Oct. 16, 2006.

U.S. Office Action from U.S. Appl. No. 10/369,148 dated Mar. 14, 2006.

U.S. Office Action from U.S. Appl. No. 10/894,165 dated Aug. 16, 2006.

U.S. Office Action from U.S. Appl. No. 12/080,594 dated Dec. 22, 2008.

U.S. Office Action from U.S. Appl. No. 11/703,358 dated Jun. 11, 2008.

Office Action from U.S. Appl. No. 11/098,997 dated August 26, 2009.

Office Action from U.S. Appl. No. 11/098,997 dated Nov. 20, 2008.

## 1

HYDROCARBON CONVERSION PROCESS  
IMPROVEMENTSCROSS-REFERENCE TO RELATED  
APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 11/098,997, filed Apr. 5, 2005, now abandoned, which is a continuation-in-part of U.S. patent application Ser. No. 10/430,240 filed Aug. 19, 2003, now U.S. Pat. No. 7,161, 10 050, which is a continuation-in-part of prior application Ser. No. 10/365,346 filed Feb. 12, 2003, now abandoned, which is a continuation of prior application Ser. No. 10/298,440 filed Nov. 19, 2002, abandoned, which is a continuation-in-part of prior application Ser. No. 10/208,068, filed Jul. 29, 2002, 15 abandoned, which is a continuation-in-part of prior application Ser. No. 10/054,004 filed Jan. 24, 2002, now U.S. Pat. No. 6,486,368, which is a continuation-in-part of prior application Ser. No. 09/951,739, filed Sep. 11, 2001, now U.S. Pat. No. 6,465,696, which is a continuation-in-part of application 20 Ser. No. 09/886,078 filed Jun. 20, 2001, now U.S. Pat. No. 6,472,572.

## CLAIM OF PRIORITY

25 Applicant claims priority based on provisional patent application Ser. No. 60/559,844, filed Apr. 6, 2004.

## TECHNICAL FIELD

This invention relates generally methods and apparatuses for synthesizing olefins, alcohols, ethers, and aldehydes from alkanes, alkenes, and aromatics, and more particularly to specific improvements in the methods and apparatuses disclosed in the patents and patent applications identified herein. 35

BACKGROUND AND SUMMARY OF THE  
INVENTION

The present application comprises a continuation-in-part 40 of application Ser. No. 10/430,240, the disclosure of which is incorporated herewith by reference as if set forth herein. The present invention comprises specific improvements in and to the methods and apparatuses disclosed and described in the patents and patent applications identified herein, specifically 45 including:

Safety Improvements;  
Use of Alternative Feedstocks;  
Process Simplification;  
Improvements to the Halogenation Step;  
Improvements to the Reproportionation Step;  
Improvements to the Solid Oxide Reaction;  
Improvements to Solid Oxide Regeneration;  
Improvements in Separations;  
Maintenance;  
Start-up;  
Shut-down;  
Materials of Construction.

## DETAILED DESCRIPTION

## Safety Improvements

1. The safety of the process may be improved by shipping 65 the halogen in solid halide form. Some specific variations include:

## 2

- a. The solid may be one of the solids that will be used in the process. One advantage is that separate halide liberation equipment may be avoided. The solid may be regenerated using air, oxygen, and/or oxygen-containing gas in the process equipment.
- b. The solid may also be a solid that is not used in the process, but rather a solid which is inexpensive, relatively non-toxic, able to liberate halide at lower temperature, liberates halide with heating rather than oxidation, less hygroscopic, less corrosive in solid form, less corrosive to regenerate, less volatile, more dense, containing a higher halogen content, disposable, and/or otherwise more convenient for shipping. Examples include:
  - i. Copper chloride, copper bromide, or copper iodide may be used for the shipment of chlorine, bromine, or iodine due to relatively low cost of copper relative to some other materials.
  - ii. Copper halides may be used for shipment for processes in which the reactive solid contains nickel, chromium, lead, cobalt, or other potentially toxic elements.
  - iii. Calcium bromide may be used to ship bromine for use in a process using a cobalt-containing metal oxide since cobalt bromide readily forms highly hydrated species.
  - iv. Copper chloride, copper bromide, or copper iodide may be used for the shipment of chlorine, bromine, or iodine since copper halides may be easily regenerated with air, oxygen, or oxygen containing gas at temperatures below those required to regenerate other solids.
  - v. Copper (II) bromide may be used to ship bromine since bromine may be liberated by heating without oxygen.
  - vi. Copper bromide may be used to ship halide for processes using iron-containing materials since iron bromide is hygroscopic, potentially volatile, and potentially corrosive.
  - vii. Pure copper halide may be used to ship halide for processes using a supported metal oxide. Such a choice will eliminate the need to transport the inert support.
  - viii. Carbon tetrabromide may be used with combustion of this material either in dedicated equipment or within a process unit, generating bromine and carbon dioxide. Such a solid is disposable, thereby eliminating the requirement of returning the solid oxide to a processing facility.
- c. The solid may also be a solid that is not used in the process so that the solid used may be shipped in oxide form, which may be more stable, less likely to sinter, dissolve, adsorb (or absorb) water (or other contaminants), or fracture.
- d. As the solid used in the process can hold substantially more bromine than the optimum level for use in the process, the solid could be shipped to an operating facility with a relatively high level of bromine that could be used to make up the small amount of bromine that may be lost during normal operation of the process.
- 2. The safety of the process may be improved by shipping the halogen in liquid halide form. The use of liquid may minimize solids handling operation and associated particulate hazards. Liquid may also be easier to handle.
  - a. Specifically, boron tribromide may be used with oxidation to boron oxide liberating bromine.
  - 3. The safety of the process may be improved by shipping the halogen in liquid alkyl halide form. The use of liquid

halides may minimize solids handling operation and associated particulate hazards. The use of liquid halides may also be easier to handle. The shipment of alkyl halide may be particularly useful for the startup of the plant and may also provide a convenient and safe way to introduce make-up halide.

4. The safety of the process may be improved by the placement of hygroscopic metal halides in selected reactors provided with a sink for water in the event of a process upset. Many metal halides are hygroscopic and will react with water to form hydrates, minimizing corrosion.

- a. The metal halides may be selected and placed so that they are molten in the hydrated form and easily removed.
- b. The metal halides may be selected based on a low melting temperature so that they can be pumped into the process in the event of an upset.

5. The bromine inventory may be reduced by utilization of the bromine separation apparatus (typically following the regeneration reactor) as the reservoir for bromine for introduction into the alkane bromination reactor or other necessary step involving bromine. This reservoir of liquid bromine will have sufficient capacity to maintain adequate pump priming and allow bromine to be pumped as a liquid rather than using more costly compressors.

6. Reactive metal oxide traps at all process vents for use in normal and emergency operations may be used to insure against release of any and all organic-bromides. These metal oxides may be regenerated to recover bromine.

7. The safety of the process may be enhanced by the use of a solid oxide to dispose of halogenated organic streams and recovery of halide by conversion to carbon dioxide, water, and solid halide. The solid oxide may be regenerated by reacting the solid halide with oxygen, liberating halogen for recycle to the process.

- a. One example is the use of CuO or CuZrO<sub>3</sub> to convert vinyl bromide to carbon dioxide, water, coke, and CuBr or CuBrZrO<sub>2</sub>.

#### Use of Alternate Feedstocks

The above-identified processes may be useful and particularly valuable with feedstocks containing otherwise difficult to separate components. The halogenation chemistry may facilitate the reactive separation of various streams including:

1. The use of streams containing alkane and olefin of the same carbon number.

- a. The olefin may be converted with molecular halogen or solid halide to the 1,2-dihalide for use as a feedstock to an epoxide process. The 1,2-dihalide will be easy to separate from the alkane. Examples include:

i. Converting the propylene in a stream containing propane and propylene to 1,2-dibromopropane and subsequently to propylene oxide. The 1,2-dibromopropane is formed by reacting the mixed hydrocarbon stream with bromine, most preferably at low temperatures where little appreciable reaction with propane occurs. Separation of propane from propylene is required in many existing plants including ethylene plants and is considered one of the most difficult separations in the chemical industry.

- b. The olefin may be converted with wet halogen to halohydrin for use as a feedstock to an epoxide process. The halohydrin will be easy to separate from the alkane. Examples include:

i. Converting the propylene in a stream containing propane and propylene to the bromohydrin and subse-

quently to propylene oxide. The propylene is converted by passing the hydrocarbon stream through bromine water.

c. The olefin may be converted with hydrogen halide to form the monohalide for use as a feedstock in an olefin, alcohol, epoxide, aldehyde, ketone, or other process. The halide will be easy to separate from the alkane. Examples include:

- i. Converting olefin in a gasoline feed to alkyl halide by reacting with hydrogen halide to form alkyl halide. The alkyl halide can be easily removed, leaving olefin-depleted gasoline.

- ii. Converting butenes to butyl halides in a mixed feed of butanes and butenes. The butyl halides may be coupled to products containing eight carbon atoms for use in gasoline.

2. The use of streams containing branched and linear alkanes resulting in product streams enriched in branched and/or linear molecules.

a. Branched alkanes containing tertiary carbon may be selectively halogenated to alkyl halide and separated, leaving a stream enriched in linear alkane. Examples include:

- i. Depletion of the branched content of detergent-range alkanes by reaction with halide, resulting in greater reactivity with the branched alkanes to branched alkyl halides. Following separation of the branched halides, the remaining stream is enriched in linear alkane. The stream rich in branched halides may be dehydrohalogenated either catalytically or using a solid oxide to create a stream rich in branched olefin for hydroformylation and conversion to branched alcohols.

b. Branched alkanes containing tertiary carbon may be selectively halogenated to alkyl halide. Following separation, the halide stream will inevitably contain some non-branched halides. By selectively dehydrohalogenating the tertiary halides, a stream containing a very high fraction of branched olefins can be separated from the remaining halides. The selective dehydrohalogenation may be conducted thermally, using a catalyst at temperatures below those required for secondary alkyl halide dehydrohalogenation, or using a solid oxide catalyst at temperatures below those required for secondary alkyl halide dehydrohalogenation. Dehydrohalogenating the remaining halides will leave a stream enriched in linear olefins.

3. The use of streams containing multiple types of branched molecules resulting in product streams enriched or depleted in molecules containing a certain type or amount of branching:

a. Streams containing linear, mono-branched, and multiply-branched alkanes may be enriched or depleted in multiply-branched product by halogenating the multiply-branched alkanes to multiply-halogenated separation. The multiply-halogenated and/or mono-halide species may be easily separated. Following the desired separation of the non-halogenated, mono-halogenated, and multi-halogenated species, and dehydrohalogenation of the halides, the various streams may be recombinined to generate the desired branching composition.

b. Streams containing branched alkanes with and without multiple branching at a single carbon (quaternary carbon) may be depleted in these quaternary carbon-containing species by halogenating the branched alkanes without the quaternary carbon, separating these halides, and dehydrohalogenating. The result will be streams

- rich in branched olefins without quaternary carbon and alkanes with quaternary carbon.
4. The use of streams containing trace amounts of impurities that are more reactive than the desired alkane reactant:
- Alkane streams containing aromatics, alcohols, olefins, aldehydes, ketones, sulfides, sulfates, or other reactive molecules may be halogenated at low temperature to selectively halogenate the impurities for removal.
  - Streams of mixed alkanes (e.g. natural gas, refinery streams) may be differentially halogenated based on differing rates of halogenation and subsequently reacted with metal oxides at lower temperatures where the non-halogenated alkanes would pass through without reaction.
  5. The use of streams of mixed alkane and olefin in coupling processes. Streams containing alkanes and olefins may be used to produce products of higher carbon number. Several process variations may be employed:

a. A process with:

- Olefin hydrohalogenation in the presence of the alkane;
  - Separation of the resulting alkyl halide from the alkane;
  - Halogenation of Alkane;
  - Separation of the resulting alkyl halide from the alkane;
  - Recycle of alkane;
  - vi. Feed of the alkyl halide to the coupling reactor.
- b. Several variations of (a) may be employed:
- Specifically, step ii (separation of alkyl halide after hydrohalogenation) may or may not be omitted.
  - The alkane may or may not be separated from the alkyl halide (step iv).
  - Reproportionation chemistry may or may not be employed.
  - The olefin and alkane may be separated at the beginning of the process.
  - Halogenation may precede hydrohalogenation, particularly if high temperature is employed to hinder addition of halogen to the olefin.
  - vi. The hydrogen halide used for hydrohalogenation may or may not be the same formed in the halogenation step.

Process Simplification

1. The halogenation and solid oxide reaction steps may be conducted in the same unit:
  - The halogenation and solid oxide reaction may be simultaneous.
  - b. The halogenation may occur first by varying the contacting of the hydrocarbon, halogen, and solid oxide.
2. The halogenation, solid oxide reaction, and solid oxide regeneration may be conducted in the same unit by introducing hydrocarbon and oxygen to a solid halide or solid halide-oxide combination. The oxygen will regenerate the solid halide generating hydrocarbon halide and solid oxide, the hydrocarbon halide will react with the oxide, generating product.

Variations include:

- Periodic switching of the direction of feed to the reactor to minimize halogen migration from the reactor.

Examples include:

- Coupling methane to heavier products by cofeeding methane and oxygen over a metal-halide-containing solid.

3. The solid oxide reaction and product separation may be conducted simultaneously when the product is lighter than the reactant the reaction is conducted in a liquid phase reactor under conditions where the product is a vapor and leaves the reactant mixture.

4. Reacting the halide-containing regeneration effluent with olefin to form dihaloalkanes to reduce the energy required for and equipment size in the halide recovery.
5. Reacting the alkane over selected metal-halides in the regeneration step to form the alkyl-halide and a metal-hydride. This would also be a safety improvement and eliminate the need for halogen separation. Materials include but are not limited to halides of boron, nickel, iron, and their mixtures as well as carbon based materials (e.g. C60).

6. Operation of the halogenation process at high halogen: alkane ratio for the feed at temperatures and pressures to maximize the production of monohalo-alkanes at 100% alkane conversion. The alkane feed may be mixed. The products which will contain multiply-halogenated species and halo acid which may be passed directly over a metal oxide bed to produce a mixture of products dependent upon the reaction conditions will be produced which will be condensed together and separated in the liquid phase by a combination of distillation and phase separation.

7. The use of a hydrogenation step to recover the over-halogenated products by reducing the halogenation to the desired degree. The use of such a step will allow for higher perpass conversion in the halogenation step. Catalysts may be used, including but not limited to Pd, Pt, Ru, Ni, Au, Cu, and their alloys.

8. Controlling the amount of hydrogen halide added to a metal oxide reactor in order to generate the heat required for an endothermic reaction.

9. The use of hydrogen halide formed in the halogenation step for conversion of byproducts or products into more useful compounds.

- a. For example, HBr could be used to hydrobrominate vinylbromide, a common undesirable by-product resulting from HBr elimination from dibromoethane back to same (or any higher vinylbromide equivalent to the corresponding dibromoalkane). In another example, HBr could be used in the acidic cleavage of ethers into alcohols and alkylbromides, the former increasing the yield of the desired alcohol product and the latter being recycled to the educt stream for reaction on the metal oxide.

50 Improvements to the Halogenation Step

1. Improvements in selectivity to desired multiply-halogenated isomers through isomerization of the multiply-halogenated species formed by halogenation. Examples include:

a. Forming dihalides dehydrohalogenating the dihalides, and rehydrohalogenating to form the desired isomers. The rehydrohalogenation may be conducted using process conditions different from the initial halogenation to enhance yield of the desired isomer. The process conditions varied may include temperature, pressure, and catalyst. Some examples include:

- i. Halogenating ethane so that it contains mixed halides including 1,1 and 1,2-dihaloethanes. Dehydrohalogenating the dihaloethanes, and rehydrohalogenating to enrich the 1,2-dihaloethane content.
- ii. Halogenating propane so that it contains mixed halides including 1,1, 2,2, 1,3, and 1,2-dihalopro-

- panes. Dehydrohalogenating the dihalopropanes, and rehydrohalogenating to enrich the 1,2-, 2,2-, 1,3, or 1,1-isomer content.
- iii. Halogenating butane so that it contains mixed dihalides. Dehydrohalogenating, and rehydrohalogenating to enrich the 2,2- or 2,3-isomer content. The 2,2- or 2,3-isomer may be reacted with a metal oxide to make methyl-ethyl ketone.
- iv. Halogenating butane so that it contains mixed tetrahalides. Dehydrohalogenating, and rehydrohalogenating to enrich the 1,2,3,4-isomer content.
- v. Halogenating cyclohexane so that it contains mixed halides including 1,1, 1,2, 1,3, and 1,4-dihaloisomers. Dehydrohalogenating, and rehydrohalogenating to enrich the 1,1, 1,2, 1,3, and 1,4-dihaloisomer content.
2. Enrichment in the primary halide content of a stream of mixed halide isomers by separating primary halides from other halide isomers. Dehydrohalogenating the other isomers, rehydrohalogenating the resulting olefins to produce a stream enriched in primary halide isomers, and returning the resulting stream to the primary halide separation step. Some variations include:
- Using selective dehydrohalogenation of the non-primary isomers to form easily separated olefin and hydrogen halide.
  - Using distillation to separate the primary and other isomers.
  - Using adsorption to separate the primary and other isomers.
  - Using a shape-selective catalyst to rehydrohalogenate the olefin, enhancing primary halide yield.
3. The use of multiple halogens to create the desired halide isomer. One halogen may be used to halogenate the hydrocarbon and be replaced by another.
4. The use a membrane reactor with halogen on one side and alkane on the other to improve selectivity to the desired halide isomer. This reactor design may improve monohalogenation, dihalogenation, and/or primary halogenation selectivity.
5. Operation of the halogenation reaction at high halogen: alkane ratio to improve conversion may result in unconverted halogen. Photoactivation of the unconverted halogen may be used at low temperature in a solid oxide bed to allow full recovery of all the halogen.

#### Improvements to the Reproportionation Step

In many processes, the overhalogenated species may be recycled to a point in the process where they are converted to the desired degree of halogenation or less than the desired degree of halogenation. The change in degree of halogenation is termed "reproportionation," and allows for the use of the carbon and hydrogen in the overhalogenated species, thus reducing feedstock loss and perhaps also allowing greater economic per-pass yield.

#### Several Improvements Include:

- A low-temperature reproportionation step, in which the halogen is redistributed among over-halogenated species, resulting in the formation of optimally halogenated species and additional very highly halogenated species.
- An example is a mixture of dibromomethane, tribromomethane, and tetrabromomethane are allowed to react, producing a stream enriched in methyl bromide and tetrabromomethane.

2. A low-temperature reproportionation step, in which the halogen is redistributed among over-halogenated species, resulting in the formation of optimally halogenated species and additional very highly halogenated species. The yield of optimally halogenated species is maximized by conducting this reproportionation under temperature, pressure, and process conditions such that the reproportionation is conducted in the liquid phase while the optimally halogenated species is predominantly in the vapor phase.
- An example is: a mixture of liquid dibromomethane, tribromomethane, and tetrabromomethane are allowed to react in the presence of a catalyst at about 30 C. As the bromine is redistributed and methyl bromide is formed, much of the methyl bromide leaves the solution and enters the vapor phase.
  - The conversion of over-halogenated hydrocarbon to carbon black or other carbon material and halogen. The carbon material may be sold and the halogen may be recycled to the process.
  - The reproportionation of overhalogenated hydrocarbon with another hydrocarbon or halohydrocarbon. Such a process may allow the recovery of the desired hydrocarbon with a loss of a less desirable material.
  - For example, dibromomethane is reacted with propane to make methyl bromide and brominated propanes. Ideally, one propane molecule can be used to convert eight dibromomethane molecules to methyl bromide. The bromine can be recovered from the brominated propane through thermal decomposition, oxidation, reaction with solid oxide, or other means.
- #### Improvements to the Solid Oxide Reaction
- A method of contacting water with alkyl halide and metal oxide in a multi-phase reactor with alkyl halide, solid oxide and optional diluent present at the bottom of the reactor with refluxing water present in a zone above the reactant mixture.
  - The product yield may be increased and process corrosivity may be reduced by conducting the solid oxide reaction in a liquid phase with water present to remove metal halide as it is formed. A specific example is:
    - The reaction is conducted in a vessel containing liquid alkyl halide, liquid water, water vapor, and solid. The water vapor condenses at the top of the reactor or is returned from an external condenser and settles through the metal oxide and alkyl halide containing phase. The water dissolves metal halide as it passes through the alkyl halide phase. The solid oxide may be supported on a plate to keep it out of the liquid water phase. The water and metal halide passes into a separate liquid phase at the bottom of the reactor where some of the water is vaporized. Variations include:
      - A batch reactor.
      - A continuous reactor in which alkyl halide, metal oxide, and water (or steam) are added continuously and metal halide solution is removed continuously to a regeneration reactor where it is dried and regenerated.
      - The use of precipitation to remove metal halide from the metal halide solution. By reducing the temperature of the solution, some of the metal halide will precipitate for regeneration. The depleted metal halide solution may be recovered by filtration, centrifugation or other solids-liquid separation methods and recycled to the reactor. Recovered solids can be dried and regenerated to metal oxide and bromine.

3. The liquid phase performance of a reactor may be improved by adding a diluent. The diluent may be, but is not limited to alkanes that are readily separated from the products and reactants.

4. The yield to desired product may be improved by introducing the stream containing hydrocarbon halide to the metal oxide in stages.

5. The yield to desired product may be improved by providing a feed of solid to a fluidized bed reactor that includes some partially or completely spent material. Spent is defined as solid with no remaining oxygen (donation) capacity or bromine capacity.

6. The yield to desired product may be improved by providing a feed of solid to a fluidized bed reactor that includes some partially coked material.

7. The solid oxide reaction may be conducted in a series of switched fixed beds, some of which are undergoing regeneration at any given time.

8. In a process for the production of olefins, the di-halogenated species may be at least partially converted to olefin using certain solids. Some examples include:

- a. The reaction of silver metal with 1,2-dibromoethane to form ethylene and silver bromide. The silver bromide may be decomposed to silver and bromine using heat or electromagnetic radiation.
- b. The reaction of copper (I) bromide with 1,2-dibromoethane to form ethylene and copper (II) bromide. The copper (II) bromide may be decomposed to copper (I) bromide and bromine using heat.
- c. The reaction of 1,2-dibromomethane with a metal oxide to form ethylene, carbon dioxide, water, and metal bromide. The metal bromide may be regenerated by reaction with oxygen.

#### Improvements to Solid Oxide Regeneration

1. Varying the temperature of solid oxide prior to oxygen introduction to change the particle size of the solid oxide to a more desirable distribution.

- a. By raising the temperature, particularly to that above the regeneration onset temperature, prior to introduction of oxygen or air, the metal oxide obtained after regeneration may be reduced in the amount of fines or agglomerates it contains.
- b. By introducing oxygen at low temperature, particularly at that below the regeneration onset temperature, the metal oxide obtained after regeneration may be reduced in the amount of fines or agglomerates it contains.

2. Increasing the temperature of solid oxide prior to oxygen introduction to dehydrogenate or desorb adsorbed hydrocarbon, reducing the amount of water and possibly carbon oxides generated in regeneration, thus reducing corrosivity and simplifying halide purification.

3. Performing a separate oxidation, particularly at low temperature, to remove adsorbed hydrocarbon reducing the amount of water and carbon oxides generated in regeneration, thus reducing corrosivity and simplifying halide purification.

4. Introducing water to the solid halide to change the particle size of the resulting solid oxide to a more desirable distribution.

- a. The water may be introduced in the gas phase.
- b. The water may be introduced in the liquid phase.
- c. The water may be introduced concurrently with or prior to the introduction of oxygen.
- d. The hydrated solid may be allowed to settle and agglomerate.
- e. The hydrated solid may be subjected to intense fluidization to break apart agglomerates.

f. A slurry or aqueous phase may be formed and dried in a manner to form the desired particle size. In particular, spray drying may be used.

5. Dissolving the active metal halide to separate it from impurities, and then converting metal halide to metal oxide.

6. The use of very high temperature regeneration to remove impurities. In particular, chlorine may be removed from metal bromide in this manner.

10 a. The combination of high temperature with heating of the solid halide prior to oxygen introduction may be particularly useful. In the case of metal bromides, this methodology may allow the removal of chlorine as  $\text{ClBr}$  or  $\text{Cl}_2$ .

15 b. The reduction of the solid halide with hydrogen or other reducing agent to remove impurities. The reduced material may be reoxidized with oxygen, air, or other oxygen containing gas.

#### Improvements in Separations

1. Separation of halogen from nitrogen, oxygen, and other non-condensables using solid adsorbents. The solid adsorbents will adsorb the halogen, which can be removed by heating the solid or reducing the pressure. The adsorbents may be, but are not limited to:

- a. Molecular sieves;
- b. Mesoporous materials;
- c. Zeolites;
- d. Silica;
- e. Alumina;
- f. Aluminosilicates;
- g. Magnesia;
- h. Activated carbon;
- i. Metal bromides;
- j. Metal oxides;

35 2. Separation of halogen from nitrogen, oxygen, and other non-condensables using reactive solid adsorbents. The solid reactive adsorbents will react with the halogen, forming a new chemical composition, from which the halogen can be removed by heating the solid or reducing the pressure, regenerating the solid. The reactive adsorbents may be, but are not limited to:

- a. Copper (I) bromide;
- b. Iron (II) bromide;
- c. Silver bromide;
- d. Carbon;
- e. Carbon, particularly fullerenes or nano-tubular carbon.
- 45 3. Removal of water from halogen by passing the mixed stream over metal halides or metal halide hydrates which may be supported or unsupported. The metal halides will form hydrates and the metal halide hydrates will form more highly hydrated species. The water can be liberated and starting material can be regenerated by heating.

50 4. Methods of removing trace amounts of halogen from product streams using reactive solids, which may or may not be regenerable. Some specific reactive solids include, but are not limited to:

- a. Copper (II) oxide;
- b. Silver;
- c. Copper;
- d. Lithium;
- e. Magnesium;
- f. Alkali metals.

55 5. Removal of residual halogen from streams by reaction with olefins. Specific examples include:

- a. Reacting with ethylene or propylene to form dihaloalkane, which can be converted to epoxide.

11

- b. Reacting with a heavy multiple-olefin to form highly halogenated species from which the halide can be recovered.
6. Removal of residual hydrocarbon halide from streams by reaction with reactive solids Some specific reactive solids include, but are not limited to:

- a. Copper (II) oxide;
- b. Silver;
- c. Copper;
- d. Lithium;
- e. Magnesium;
- f. Alkali metals.

7. Separation of primary, secondary, and/or tertiary alkyl halides by selective dehydrohalogenation of selected species, separation of the olefin and hydrogen halide from the remaining alkyl halide, and recombination of the hydrogen halide and olefin to form alkyl halides. Examples include, but are not limited to:

- a. Separation of primary from secondary alkyl halides by selectively dehydrohalogenating the secondary alkyl halides to olefin and hydrogen halide, separation of the olefin and hydrogen halide from the primary halide, and recombination of the hydrogen halide and olefin to form secondary and possibly primary halide.
- b. Separation of primary and secondary alkyl halides from tertiary alkyl halides by selectively dehydrohalogenating the tertiary alkyl halides to olefin and hydrogen halide, separation of the olefin and hydrogen halide from the primary and secondary alkyl halide, and recombination of the hydrogen halide and olefin to form alkyl halide.

8. Removal of sulfur-containing compounds from a hydrocarbon feed by reacting the feed with dry halogen to form sulfur, which can be removed as a solid from the sulfur-depleted hydrocarbon and hydrocarbon halide stream.

9. Removal of sulfur-containing compounds from a hydrocarbon feed by reacting the feed with dry halogen to form sulfur, which can be removed as a solid from the sulfur-depleted hydrocarbon and hydrocarbon halide stream.

10. Removal of carbon dioxide from a stream by reacting with a carbonate-forming material such as calcium oxide. The carbonate may be used in a hydrogen-halide recovery section of the plant. The carbonate will react with hydrogen halide, liberating water and carbon dioxide and producing solid halide, which can be regenerated and recycled to the carbon dioxide separation section.

11. Removal of arsenic, mercury, heavy metal-containing compounds from a hydrocarbon feed by reacting the feed with dry halogen to form solid metal compounds, solid metal halides or halogenated metal hydrocarbons, which can be easily separated.

12. Removing adsorbed product from the solid by rinsing with a compound that is easily separated from the product. Such a rinsing agent may be pentane or other alkane.

13. Removing adsorbed product from the solid by steam distillation.

#### Maintenance

1. A method of removing coke from reactors by reacting with bromine to form volatile carbon bromides. The carbon bromides may be used in the process in a reportionation step, thus producing product from the coke.
2. A method of removing coke from reactors by reacting with hydrogen bromide to form volatile hydrocarbon bromides. The carbon bromides may be used in the process in a reportionation step, thus producing product from the coke.

12

#### Start-Up

1. Starting the process with some or all of the solid in the halide or partially halogenated form may provide a number of benefits including:

- a. The ability to start the regeneration reactor early in the start up sequence.
- b. Reduction in the amount of oxygen carried over into the bromine separation unit.
- c. Reduction in the heat generated in the metal oxide reactor or hydrogen halide neutralization step.
- d. Reduction in the amount of adsorbed hydrocarbon and thus reduction in the amount of water and carbon dioxide generated in the regeneration unit.
- e. Reduction in unfavorable changes in the particle size distribution of the metal oxide.
- f. Improvement in the packing of a fixed bed reactor.
- g. Providing a hygroscopic metal halide to reactively remove water during upsets.

2. Starting the process with some or all of the solid in the oxide or partially oxygenated form may provide a number of benefits including:

- a. The ability to start the solid oxide reactor early in the start up sequence.
- b. Reduction in the amount of bromine generated in the regeneration reactor.
- c. Reduction in the heat generated in the metal oxide reactor or hydrogen halide neutralization step.
- d. Reduction in the amount of adsorbed hydrocarbon and thus reduction in the amount of water and carbon dioxide generated in the regeneration unit.
- e. Reduction in unfavorable changes in the particle size distribution of the metal
- f. Improvement in the packing of a fixed bed reactor.
- g. Providing a hygroscopic metal halide to reactively remove water during upsets.

3. Starting the process with a solid which has undergone a number of regeneration cycles may offer benefits including:

- a. Reduction in byproducts.
- b. Improved chemical and thermal stability of reactors
- 4. Starting the process with the halogen present in part or completely as alkyl halide may be desirable for a number of reasons including:

- a. Less free halogen present during start-up.
- b. Ability to start the halide separations section early in the start-up sequence with no hydrogen halide or water present.

#### Shut-Down

1. Stopping the process with the halide in metal halide and/or alkyl halide form may improve safety, reduce corrosion, and improve maintenance accessibility.

2. Introducing reactive components into certain sections of the plant may provide a sink for halogen or hydrogen halide, improving safety, reducing corrosion and improving accessibility. An example of such a component is olefin.

#### Materials of Construction

1. The reactors for alkane halogenation and metathesis consisting of materials to minimize corrosion including but not limited to:

- a. Stainless steel;
- b. Silicon carbide;
- c. Glass lined steel;
- d. Titanium;
- e. Carbon fiber.

## 13

2. Process components operating at temperatures below 300 C. constructed from:

- a. Teflon;
- b. Glass.

What is claimed is:

1. A method for use in conjunction with a process of hydrocarbon conversion comprising:  
 providing a first stream comprising a halide;  
 contacting the first stream with a solid material, wherein the solid material retains at least a portion of the halide in the first stream;  
 removing at least a portion of the halide retained by the solid material from the solid material; and  
 contacting at least a portion of the halide removed from the solid material with a hydrocarbon feed, and  
 reacting at some hydrocarbons in the hydrocarbon feed with at least a portion of the halide removed from the solid material to form a second stream comprising an alkyl halide.

2. The method of claim 1 wherein the solid material comprises a reactive adsorbent.

3. The method of claim 2 wherein the reactive adsorbent comprises at least one material selected from the group consisting of: copper (I) bromide, iron (II) bromide, silver bromide, carbon, carbon fullerenes and nano-tubular carbon.

4. The method of claim 1 wherein the solid material is a reactive solid, wherein the reactive solid comprises at least one material selected from the group consisting of: copper (II) oxide, silver, copper, lithium, magnesium, and an alkylide metal.

5. The method of claim 1 wherein the hydrocarbon feed comprises at least one hydrocarbon selected from the group consisting of: methane, ethane, propane, butane, isobutene, pentane, hexane, and cyclohexane.

6. The method of claim 1 further comprising:  
 separating any higher alkyl halides from the second stream, wherein the higher alkyl halides comprise any alkyl halide with a plurality of halogen atoms;  
 reproportionating the higher alkyl halides with at least some of the hydrocarbon feed to form at least some alkyl monohalide; and  
 combining at least some of the alkyl monohalide with the second stream.

7. The method of claim 1 wherein the solid material comprises an adsorbent, wherein the adsorbent comprises at least one structure selected from the group consisting of: a molecular sieve, a mesoporous material, a zeolite, a silica, an alumina, an aluminosilicate, a magnesia, an activated carbon, a metal bromide, and a metal oxide.

8. A method for use in conjunction with a process of hydrocarbon conversion comprising:

providing a first stream comprising a halide;  
 contacting the first stream with a solid material, wherein the solid material retains at least a portion of the halide in the first stream;  
 removing at least a portion of the halide retained by the solid material;  
 reacting at least a portion of the halide removed from the solid material with a hydrocarbon feed to form a second stream, wherein the second stream comprises an alkyl halide; and  
 contacting the second stream with a catalyst to cause a coupling reaction, wherein the coupling reaction creates at least a product stream.

9. The method of claim 8 wherein the solid material comprises an adsorbent, wherein the adsorbent comprises at least one structure selected from the group consisting of: a molecular sieve, a mesoporous material, a zeolite, a silica, an alumina, an aluminosilicate, a magnesia, an activated carbon, a metal bromide, and a metal oxide.

## 14

10. The method of claim 8 wherein the solid material comprises a reactive adsorbent, wherein the reactive adsorbent comprises at least one material selected from the group consisting of: copper (I) bromide, iron (II) bromide, silver bromide, carbon, carbon fullerenes and nano-tubular carbon.

11. The method of claim 8 wherein the solid material is a reactive solid, wherein the reactive solid comprises at least one material selected from the group consisting of: copper (II) oxide, silver, copper, lithium, magnesium, and an alkylide metal.

12. The method of claim 8 wherein the catalyst comprises at least one material selected from the group consisting of: a metal oxide, a metal oxyhydrate, a hydrate of a halide, a hydrate of a sulfide, a hydrate of a carbonate, a hydrate of a phosphate, a hydrate of a phosphide, a hydrate of a nitride, and a hydrate of a nitrate.

13. The method of claim 8 wherein the second stream is contacted with the catalyst in stages.

14. The method of claim 8 wherein the second stream is contacted with the catalyst in a plurality of switched fixed beds.

15. A method for use in conjunction with a process of hydrocarbon conversion comprising:

providing a first stream comprising a halide;  
 contacting the first stream with a solid material, wherein the solid material retains at least a portion of the halide in the first stream;  
 removing at least a portion of the halide retained by the solid material;  
 reacting at least a portion of the halide removed from the solid material with a hydrocarbon in a feed comprising hydrocarbons to form a second stream, wherein the second stream comprises an alkyl halide;  
 contacting the second stream with a metal oxide to cause a coupling reaction, wherein the coupling reaction produces a metal halide and a product comprising at least one component selected from the group consisting of: an olefin, an alcohol, an ether, and an aldehyde; and  
 contacting the metal halide with an oxidizer to produce at least some of the metal oxide and at least some regenerated halide.

16. The method of claim 15 wherein the solid material comprises an adsorbent, wherein the adsorbent comprises at least one structure selected from the group consisting of: a molecular sieve, a mesoporous material, a zeolite, a silica, an alumina, an aluminosilicate, a magnesia, an activated carbon, a metal bromide, and a metal oxide.

17. The method of claim 15 wherein the solid material comprises a reactive adsorbent, wherein the reactive adsorbent comprises at least one material selected from the group consisting of: copper (I) bromide, iron (II) bromide, silver bromide, carbon, carbon fullerenes and nano-tubular carbon.

18. The method of claim 15 wherein the solid material is a reactive solid, wherein the reactive solid comprises at least one material selected from the group consisting of: copper (II) oxide, silver, copper, lithium, magnesium, and an alkylide metal.

19. The method of claim 15 further comprising:  
 separating any higher alkyl halides from the second stream, wherein the higher alkyl halides comprise any alkyl halide with a plurality of halogen atoms;  
 reproportionating the higher alkyl halides with at least some of the hydrocarbon feed to form at least some alkyl monohalide; and  
 combining at least some of the alkyl monohalide with the second stream.

20. The method of claim 15 wherein the regenerated halide forms at least a portion of the first stream.